

A New Expression for the Form Stress Term in the Vertically Lagrangian Mean Framework for the Effect of Surface Waves on the Upper-Ocean Circulation

HIDENORI AIKI

Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

RICHARD J. GREATBATCH

GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, Kiel, Germany

(Manuscript received 22 November 2012, in final form 27 February 2013)

ABSTRACT

There is an ongoing discussion in the community concerning the wave-averaged momentum equations in the hybrid vertically Lagrangian and horizontally Eulerian (VL) framework and, in particular, the form stress term (representing the residual effect of pressure perturbations) that is thought to restrict the handling of higher-order waves in terms of a perturbation expansion. The present study shows that the traditional pressure-based form stress term can be transformed into a set of terms that do not contain any pressure quantities but do contain the time derivative of a wave-induced velocity. This wave-induced velocity is referred to as the pseudomomentum in the VL framework, as it is analogous to the generalized pseudomomentum in Andrews and McIntyre. This enables the second expression for the wave-averaged momentum equations in the VL framework (this time for the development of the total transport velocity minus the VL pseudomomentum) to be derived together with the vortex force. The velocity-based expression of the form stress term also contains the residual effect of the turbulent viscosity, which is useful for understanding the dissipation of wave energy leading to a transfer of momentum from waves to circulation. It is found that the concept of the virtual wave stress of Longuet-Higgins is applicable to quite general situations: it does not matter whether there is wind forcing or not, the waves can have slow variations, and the viscosity coefficient can vary in the vertical. These results provide a basis for revisiting the surface boundary condition used in numerical circulation models.

1. Introduction

Wave-averaged momentum equations for the effect of surface gravity waves on circulation in the upper ocean have been derived in either the three-dimensional Eulerian mean framework (e.g., McWilliams et al. 2004; Lane et al. 2007) or the three-dimensional Lagrangian mean framework (e.g., Pierson 1962; Andrews and McIntyre 1978, hereafter AM78; Ardhuin et al. 2008b). An ongoing discussion in recent studies is whether the corresponding equations can be obtained from the vertically Lagrangian and horizontally Eulerian (VL)-mean framework that was introduced in prototype form by Mellor (2003) and Broström et al. (2008). Despite the

discussion by Lane et al. (2007) and Ardhuin et al. (2008a), how to derive the wave-averaged momentum equations with the so-called vortex force “using the VL framework” remains an open question in the oceanographic community. On the other hand, as noted by Aiki and Greatbatch (2012, hereafter AG12), the VL framework offers a concise treatment of the viscous boundary condition at the sea surface, because the viscosity term of the wave-averaged momentum equations in the VL framework is written in a flux-divergence form. Thus, as shown in AG12, the VL framework can be used as an alternative to the three-dimensional Lagrangian framework of Pierson (1962) for explaining the virtual wave stress (VWS) of Longuet-Higgins (1953, 1960) [not to be confused with the radiation stress of Longuet-Higgins and Stewart (1964)]. To our knowledge the VWS has not been explained using the three-dimensional Lagrangian framework of AM78 in previous studies, apart from the attempt by Ardhuin et al. (2008b). The goal of the present study is to clarify the relationship between

Corresponding author address: Hidenori Aiki, Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology, 3173-25, Showamachi, Kanazawaku, Yokohama 236-0001, Japan.
E-mail: aiki@jamstec.go.jp

the VL framework and the three-dimensional Lagrangian framework. We show that the traditional pressure-based expression of the form stress term in the VL framework can be rewritten as a velocity-based expression, which we argue is the cornerstone for settling the discussion regarding both the vortex force in the VL framework and the relevance of the VWS to quite general situations.

The work of Lagrange (1788), who developed two different expressions for the momentum equations in Lagrangian coordinates, is highly relevant to the present study. Let (x^e, y^e, z^e) be the instantaneous position of a fluid particle in the Eulerian–Cartesian coordinates and $(u, v, w) \equiv (dx^e/dt, dy^e/dt, dz^e/dt)$ be velocity where d/dt is the material derivative operator. Each fluid particle can be labeled by either its initial position (Lagrange 1788; Lamb 1932) or its low-pass-filtered position (AM78) and is here symbolized as (a, b, c) . The first expression (hereafter referred to as the direct expression) for the momentum equations in the three-dimensional Lagrangian coordinates reads

$$\begin{pmatrix} \rho du/dt - Q^u \\ \rho dv/dt - Q^v \\ \rho dw/dt - Q^w \end{pmatrix} = - \begin{pmatrix} x_a^e & y_a^e & z_a^e \\ x_b^e & y_b^e & z_b^e \\ x_c^e & y_c^e & z_c^e \end{pmatrix}^{-1} \begin{pmatrix} p_a \\ p_b \\ p_c \end{pmatrix}, \quad (1)$$

where ρ is density and (Q^u, Q^v, Q^w) is the sum of the effects of viscosity, gravitational acceleration, and the rotation of the earth. To obtain an expression in the three-dimensional Lagrangian coordinates, the pressure gradient in the Eulerian coordinates $(p_{x^e}, p_{y^e}, p_{z^e})$ has been rewritten using the pressure gradient in the three-dimensional Lagrangian coordinates (p_a, p_b, p_c) based on the standard chain rule between partial differentials [see (C) on p. 445 of Lagrange (1788)].¹ The Lagrangian average of (1) yields development equations for the Lagrangian mean (LM) velocity, which is the sum of the Eulerian mean (EM) velocity and the Stokes-drift velocity (Stokes 1847). Although the pressure gradient term of (1) is complicated, the forcing term is simpler than that in the second expression shown below, so that the direct expression has been often used in the studies of viscous surface waves [Chang 1969; Ünlüata and Mei 1970; Weber 1983; Jenkins 1987; Piedra-Cueva 1995; Ng 2004; all of these studies used (9) of Pierson (1962), see our Table 1].

The second expression (hereafter referred to as the transformed expression) for the momentum equations

in the three-dimensional Lagrangian coordinates is obtained by multiplying (1) with the coordinate transformation matrix associated with the chain rule to yield [see (D) on p. 446 of Lagrange (1788)]

$$\begin{pmatrix} x_a^e & y_a^e & z_a^e \\ x_b^e & y_b^e & z_b^e \\ x_c^e & y_c^e & z_c^e \end{pmatrix} \begin{pmatrix} \rho du/dt - Q^u \\ \rho dv/dt - Q^v \\ \rho dw/dt - Q^w \end{pmatrix} = - \begin{pmatrix} p_a \\ p_b \\ p_c \end{pmatrix}. \quad (2)$$

Although the lhs is complicated, AM78 have presented a straightforward manipulation to render the Lagrangian average of (2) into development equations for the difference between the LM velocity and the generalized pseudomomentum (or wave momentum, see section 3 in AM78).² Using this result of AM78, Leibovich (1980) has presented wave-averaged momentum equations that include the so-called vortex force

$$(\partial_{t^e} + \bar{\mathbf{V}}^e \cdot \nabla^e + \bar{w}^e \partial_{z^e}) \bar{\mathbf{V}}^e = -\nabla^e \mathcal{P} + \mathbf{V}^{\text{Stokes}} \times (\nabla^e \times \bar{\mathbf{V}}^e) \quad (3a)$$

and

$$(\partial_{t^e} + \bar{\mathbf{V}}^e \cdot \nabla^e + \bar{w}^e \partial_{z^e}) \bar{w}^e = -\partial_{z^e} \mathcal{P} - g + \mathbf{V}^{\text{Stokes}} \cdot (\partial_{z^e} \bar{\mathbf{V}}^e - \nabla^e \bar{w}^e), \quad (3b)$$

where $\bar{\mathbf{V}}^e$ and \bar{w}^e are the horizontal and vertical components of the EM velocity, respectively; $\mathbf{V}^{\text{Stokes}}$ is the horizontal component of the Stokes-drift velocity associated with surface gravity waves; $\partial_{t^e}, \nabla^e = (\partial_{x^e}, \partial_{y^e})$, and ∂_{z^e} are the temporal, horizontal, and vertical gradient operators, respectively, in the Eulerian–Cartesian coordinates (see Table 2); g is gravitational acceleration; and \mathcal{P} symbolizes the sum of the Lagrangian average of nonhydrostatic pressure p and other scalar quantities (associated with the so-called Bernoulli head). In (3a)–(3b), the Coriolis term and the viscosity term are omitted for simplicity. The last term of each of (3a)–(3b) is the vortex force. The vortex force represents the interaction between an EM shear flow and the Stokes-drift flow associated with surface waves and is appropriate

¹The chain rule is $\begin{pmatrix} \partial_a \\ \partial_b \\ \partial_c \end{pmatrix} = \begin{pmatrix} x_a^e & y_a^e & z_a^e \\ x_b^e & y_b^e & z_b^e \\ x_c^e & y_c^e & z_c^e \end{pmatrix} \begin{pmatrix} \partial_{x^e} \\ \partial_{y^e} \\ \partial_{z^e} \end{pmatrix}$.

²For irrotational wave motions in the vertical plane, the analytical expression of the generalized pseudomomentum of AM78 is identical to that of the Stokes-drift velocity at the leading order in terms of an asymptotic expansion, with a consequence that “the LM velocity minus the generalized pseudomomentum” approximates to the EM velocity. This is why (3a)–(3b) have been written for the development of the EM velocity. The LM velocity minus the generalized pseudomomentum has been referred to as the quasi-EM velocity in Jenkins (1989) and Ardhuin et al. (2008a,b).

TABLE 1. Classification of previous studies based on the form of the Lagrangian momentum equation that is used.

Direct expression	Transformed expression
Lagrange (1788), (C) on p. 445	Lagrange (1788), (D) on p. 446
Pierson (1962), (5) and (9)	Lamb (1932), second equation on p. 13
Andrews and McIntyre (1978), (8.7a)	Pierson (1962), (4) and (10)
Mellor (2003) and Aiki and Greatbatch (2012)	Andrews and McIntyre (1978), (3.8)
	This study

to describe the maintenance of Langmuir circulations (LCs; Langmuir 1938). LCs play an important role in the vertical mixing of the surface mixed layer of the ocean (e.g., Skillingstad and Denbo 1995; Polton and Belcher 2007; Kukulka et al. 2010). Some prototypes of the vortex force have been derived by Craik and Leibovich (1976, hereafter CL76) using EM vorticity equations and by Garrett (1976) using EM depth-integrated momentum equations. Besides the theory of LCs, the vortex force has also been used in the modeling of the circulation in an inner-coastal shelf region (e.g.,

McWilliams et al. 2004; Tang et al. 2007; Arduin et al. 2008b).

It can be said that the thickness-weighted-mean (TWM) momentum equations of Mellor (2003), Broström et al. (2008), and AG12 correspond to the Lagrangian average of (1), which is the direct expression of the Lagrangian momentum equations. This is because (i) the TWM momentum equations are written for the development of the TWM velocity whose role corresponds to that of the LM velocity in the three-dimensional Lagrangian framework and (ii) the TWM momentum

TABLE 2. List of symbols, where A is an arbitrary quantity. Variables (x^e, y^e, z^e, t^e) and ϖ are the same as (x^c, y^c, z^c, t^c) and w^* in Jacobson and Aiki (2006), AG12, and AG13.

(x^e, y^e, z^e)	Eulerian–Cartesian coordinates
(a, b, c)	Three-dimensional Lagrangian coordinates
(x, y, z)	VL coordinates
\bar{A}	Time mean in the Eulerian–Cartesian coordinates
$\hat{A} \equiv \overline{z_z^e A}$	TWM in the VL coordinates
\bar{A}	Unweighted time mean in the VL coordinates
$A' \equiv A - \bar{A}$	Deviation from the Eulerian mean, compared at fixed z^e ($\bar{A}' = 0$)
$A'' \equiv A - \hat{A}$	Deviation from the TWM, compared at fixed z ($\overline{z_z^e A''} = 0$)
$A''' \equiv A - \bar{A}$	Deviation from the unweighted mean, compared at fixed z ($\overline{A'''} = 0$)
$\mathbf{V} \equiv (\partial_x, \partial_y)$	Lateral gradient in the VL coordinates ($\mathbf{V}z = 0, \mathbf{V}z^e = z'''$)
$\nabla^e \equiv (\partial_{x^e}, \partial_{y^e})$	Horizontal gradient in Eulerian–Cartesian coordinates [$\nabla^e = \mathbf{V} - (\mathbf{V}z^e)\partial_{z^e}$]
$\mathbf{V} \equiv (u, v)$	Horizontal component of velocity
w	Vertical component of velocity
$\varpi \equiv (w - z_z^e - \mathbf{V} \cdot \nabla z^e)/z_z^e$	Vertical velocity associated with volume flux through surface of fixed z
$(\hat{\mathbf{V}}, \hat{w})$	TWM velocity
$(\hat{\mathbf{V}}, \hat{\varpi})$	Total transport velocity ($\mathbf{V} \cdot \hat{\mathbf{V}} + \hat{\varpi}z = 0$)
$(\mathbf{V}^{\text{qs}}, w^{\text{qs}}) \equiv (\hat{\mathbf{V}} - \overline{\mathbf{V}^e}, \hat{\varpi} - \overline{w^e})$	Quasi-Stokes velocity ($\mathbf{V} \cdot \mathbf{V}^{\text{qs}} + w^{\text{qs}}z = 0$)
η	Sea surface height
p	Sum of oceanic nonhydrostatic pressure and atmospheric sea surface pressure
$\mathcal{F}\mathcal{S}^{\mathbf{V}}$	Divergence of form stress $\equiv -[z'''\overline{\mathbf{V}(p'' + \eta''')}]_z + \mathbf{V}(z'''\overline{p_z''})$
$\mathcal{R}\mathcal{S}^A$ for $A = u, v,$ and w	Divergence of Reynolds stress $\equiv \mathbf{V} \cdot (z_z^e \mathbf{V}'' A'') + (z_z^e \overline{\varpi'' A''})_z$
$\mathcal{D}_t \equiv \partial_t + \mathbf{V} \cdot \mathbf{V} + \varpi \partial_z$	Introduced in (11a)–(11b)
$\hat{\mathcal{D}}_t \equiv \partial_t + \hat{\mathbf{V}} \cdot \mathbf{V} + \hat{\varpi} \partial_z$	Introduced in (9a)–(9b)
$\hat{\mathcal{D}}_T \equiv \partial_T + \hat{\mathbf{V}}_2 \cdot \mathbf{V} + \hat{\varpi}_2 \partial_z$	Introduced in (23b)–(23c)
$\overline{\mathcal{D}}_T^e \equiv \partial_T + \overline{\mathbf{V}}_2 \cdot \mathbf{V} + \overline{w}_2^e \partial_z$	Introduced in (B2)
\mathcal{A}	Amplitude of $O(\alpha)$ wave
α	Surface slope of $O(\alpha)$ wave
$\kappa \equiv \sqrt{k^2 + l^2}$	Horizontal wavenumber of $O(\alpha)$ wave
σ	Frequency of $O(\alpha)$ wave
$\theta \equiv kx + ly - \sigma\tau$	Phase of $O(\alpha)$ wave
∂_τ	Time derivative operator for wave quantities
∂_T	Time derivative operator for mean quantities ($\partial_t = \partial_\tau + \alpha'' \partial_T$)
$\hat{\mathbf{V}}$	Lateral gradient for wave quantities
$\overline{\mathbf{V}}$	Lateral gradient for mean quantities ($\mathbf{V} = \hat{\mathbf{V}} + \alpha'' \overline{\mathbf{V}}$)

equations contain the form stress term representing the residual effect of pressure perturbations. Nevertheless, in contrast to the Lagrangian average of (1), the TWM momentum equations contain the horizontal Reynolds stress term, which originates from the fact that the VL framework is Eulerian in the horizontal direction.

No previous study has attempted to derive the transformed expression for the TWM momentum equations corresponding to the Lagrangian average of (2). This is why, so far, the vortex force has not been obtained from the family of equations in the VL framework despite the discussion relating to this issue in the literature (cf. Jenkins and Arduin 2004; Lane et al. 2007; Arduin et al. 2008a; Broström et al. 2008). A key step is the derivation of a new expression for the form stress written entirely using velocity variables, the subject of section 2 of the present study. In section 3 we show that the velocity-based expression of the form stress term contains the time derivative of a wave-induced velocity, resulting in the transformed expression for the TWM momentum equations, namely the expression rewritten for the development of the EM velocity, that contain the vortex force. We then introduce turbulent viscosity in section 4 to show that the velocity-based expression of the form stress term also contains the residual effect of viscosity and is useful for understanding the issue of how the dissipation of wave energy leads to the transfer of momentum from waves to circulation. Our finding is that the VWS of Longuet-Higgins (1953, 1960) is applicable in quite general situations and needs to be taken into account when considering the boundary conditions used in numerical circulation models. Section 5 presents a brief summary.

2. Mathematical development

We consider incompressible inviscid water of constant, uniform density in a nonrotating frame (rotation and viscosity are introduced in section 4).

a. The thickness-weighted-mean equations of Aiki and Greatbatch (2012)

The incompressible condition and the momentum equations in the VL coordinates (x, y, z, t) of AG12 are

$$(z_z^e)_t + \mathbf{V} \cdot (z_z^e \mathbf{V}) + (z_z^e \varpi)_z = 0, \quad (4a)$$

$$(\partial_t + \mathbf{V} \cdot \nabla + \varpi \partial_z) z^e = w, \quad (4b)$$

$$(\partial_t + \mathbf{V} \cdot \nabla + \varpi \partial_z) \mathbf{V} = -\nabla(p + \eta) + p_{z^e} \nabla z^e, \quad \text{and} \quad (4c)$$

$$(\partial_t + \mathbf{V} \cdot \nabla + \varpi \partial_z) w = -p_{z^e}, \quad (4d)$$

where $z^e = z^e(x, y, z, t)$ is the instantaneous height of fluid particles in the standard Eulerian–Cartesian coordinates. The symbol ε , rather than c as used in AG12 and Aiki and Greatbatch (2013, hereafter AG13), is used in the present study to preserve consistency with (1) and (2). In addition, the symbol w^* in AG12 and AG13 has been replaced by the new symbol ϖ in the present study for convenience. The vertical coordinate $z \equiv \bar{z}^e$ is a low-pass-filtered height coordinate and z_z^e is the thickness. The horizontal coordinates x and y are the same as the Eulerian–Cartesian coordinates. The quantity $\mathbf{V} = (u, v)$ is the horizontal velocity vector, w is the vertical component of velocity, ϖ represents water flux through the surfaces of fixed z , $\nabla \equiv (\partial_x, \partial_y)$ is the lateral gradient operator along the surfaces of fixed z , and $\nabla z = 0$ is understood. The quantity p is the sum of the oceanic nonhydrostatic and atmospheric pressure, and η is the instantaneous sea surface height. Table 2 presents a list of the symbols used in the text. All variables and quantities (such as $x, y, z, t, u, v, w, \varpi, z^e, p$, and η) in the present manuscript have been non-dimensionalized, as in appendix A of AG13. The non-dimensionalization is not essential but serves to simplify the mathematics.

The difference between the three-dimensional Lagrangian coordinates of AM78 and the VL coordinates is illustrated in Fig. 1. As the wave propagates rightward, the control cell of the three-dimensional Lagrangian coordinates (blue) rotates clockwise and returns to its original position. The movement of the control cell captures only high-frequency fluid motion (rather than the full motion of each fluid particle), which is why the control cell does not drift away despite the Stokes drift and even though there could also be a background EM flow present in the horizontal and vertical directions. The control cell of the VL coordinates (red) moves like a piston whose thickness stretches and shrinks.

Momentum equations in a flux-divergence form can be obtained by multiplying each of (4c) and (4d) by the thickness z_z^e and then using (4a) to give

$$(z_z^e \mathbf{V})_t + \nabla \cdot (z_z^e \mathbf{V} \mathbf{V}) + (z_z^e \varpi \mathbf{V})_z = -z_z^e \nabla(p + \eta) + p_z \nabla z^e \quad \text{and} \quad (5a)$$

$$(z_z^e w)_t + \nabla \cdot (z_z^e \mathbf{V} w) + (z_z^e \varpi w)_z = -p_z, \quad (5b)$$

where $z_z^e p_{z^e} = p_z$ is understood. Low-pass temporal filtering each of (4a), (5a), and (5b) yields TWM equations for incompressibility and the horizontal and vertical components of momentum




FIG. 1. Illustration of the phase cycle of a wave propagating in the direction of the x^ε axis. A control volume element in (a) the generalized-Lagrangian-mean coordinates of AM78 and (b) the VL coordinates of the present study is shaded in blue and red, respectively, with its low-pass-filtered height, as measured in each coordinate system, indicated by horizontal lines, and the reference horizontal position indicated by vertical lines. Each color line indicates a material surface that is formed by connecting the instantaneous position of water particles whose three-dimensional Lagrangian low-pass-filtered height is a given value. Adapted from AG13.

$$\nabla \cdot \hat{\mathbf{V}} + \hat{\omega}_z = 0, \quad (6a)$$

$$\hat{\mathbf{V}}_t + \nabla \cdot (\hat{\mathbf{V}}\hat{\mathbf{V}}) + (\hat{\omega}\hat{\mathbf{V}})_z + \mathcal{RS}^{\mathbf{V}} = -\nabla(\bar{p} + \bar{\eta}) + \mathcal{FS}^{\mathbf{V}},$$

and

$$(6b)$$

$$\hat{w}_t + \nabla \cdot (\hat{\mathbf{V}}\hat{w}) + (\hat{\omega}\hat{w})_z + \mathcal{RS}^w = -\bar{p}_z, \quad (6c)$$

where $\bar{z}_z^\varepsilon = 1$ (i.e., $\bar{z}^\varepsilon = z$) has been used.³ The hat symbol is the TWM operator ($\hat{A} \equiv \overline{z_z^\varepsilon A}$ for an arbitrary

quantity A). The symbols $\mathcal{RS}^{\mathbf{V}}$ and \mathcal{RS}^w in (6b)–(6c) are the Reynolds stress terms defined by

$$\mathcal{RS}^A \equiv \nabla \cdot (\overline{z_z^\varepsilon \mathbf{V}'' A''}) + (\overline{z_z^\varepsilon \varpi'' A''})_z, \quad (7)$$

for $A = u, v$, and w . The double-prime symbol is the deviation from the TWM ($A'' = A - \hat{A}$, compared at fixed z). The last term of (7) is given by ϖ'' (not w'') and thus is nearly zero. The fact that ϖ'' is nearly zero is attributed to the way the VL coordinates have been designed so that ϖ is associated with low-frequency fluid motions and not with the waves themselves. In particular AG12 have shown that $\varpi'' = \mathbf{V}'' \cdot \nabla \bar{\eta}$ at the sea surface, which implies that the Reynolds stress vector $(\overline{z_z^\varepsilon \mathbf{V}'' A''}, \overline{z_z^\varepsilon \varpi'' A''})$ is aligned along the mean slope of the sea surface. The symbol $\mathcal{FS}^{\mathbf{V}}$ in (6b) is the form stress term defined by

³The TWM equations originate from studies on mesoscale eddies (cf. Gallimore and Johnson 1981; Andrews 1983; Bleck 1985; de Szoeke and Bennett 1993; Greatbatch 1998; Iwasaki 2001; Greatbatch and McDougall 2003; Aiki and Richards 2008; Young 2012).

$$\begin{aligned}\mathcal{F}S^{\mathbf{V}} &\equiv -\overline{z_z'''\nabla(p'''+\eta''')} + \overline{p_z'''\nabla z'''} \\ &= -[z_z'''\nabla(p'''+\eta''')]_z + \nabla(\overline{z_z'''}p_z'''),\end{aligned}\quad (8)$$

where the triple-prime symbol is the deviation from the unweighted mean ($A''' \equiv A - \overline{A}$, compared at fixed z). It should be noted that $z_z''' \equiv z^e - \overline{z^e} = z^e - z$.

For low-pass-filtered quantities, the VL coordinates (x, y, z, t) correspond to the standard Eulerian–Cartesian coordinates (Jacobson and Aiki 2006). A nice feature of the total transport velocity $(\hat{\mathbf{V}}, \hat{\omega})$ in (6a)–(6c) is that both the incompressible condition (6a) and the kinematic boundary condition ($\hat{\omega} = \overline{\eta}_t + \hat{\mathbf{V}} \cdot \nabla \overline{\eta}$ at the sea surface, see AG12) are always satisfied without relying on an asymptotic expansion approach. The incompressibility condition allows us to rewrite the TWM momentum equations (6b)–(6c) in the form

$$\hat{D}_t \hat{\mathbf{V}} + \mathcal{R}S^{\mathbf{V}} = -\nabla(\overline{p} + \overline{\eta}) + \mathcal{F}S^{\mathbf{V}} \quad \text{and} \quad (9a)$$

$$\hat{D}_t \hat{\omega} + \mathcal{R}S^{\omega} = -\overline{p}_z, \quad (9b)$$

where $\hat{D}_t \equiv \partial_t + \hat{\mathbf{V}} \cdot \nabla + \hat{\omega} \partial_z$ is the material derivative operator based on the total transport velocity. The quantities $\hat{\omega}$ and $\hat{\omega}$ are not the same mathematically but the difference is negligible as far as the present study is concerned, as is demonstrated in section 3.

The wave-induced velocity in Mellor (2003) can be called the quasi-Stokes velocity following McDougall and McIntosh (2001). The quasi-Stokes velocity ($\mathbf{V}^{\text{qs}}, \omega^{\text{qs}}$) refers to the difference between the total transport velocity $(\hat{\mathbf{V}}, \hat{\omega})$ and the EM velocity $(\overline{\mathbf{V}}^e, \overline{\omega}^e)$

$$\mathbf{V}^{\text{qs}} \equiv \hat{\mathbf{V}} - \overline{\mathbf{V}}^e (= \overline{\mathbf{V}} + \overline{z_z'''\mathbf{V}'''} - \overline{\mathbf{V}}^e) \quad \text{and} \quad (10a)$$

$$\omega^{\text{qs}} \equiv \hat{\omega} - \overline{\omega}^e (= \overline{\omega} - \overline{\mathbf{V}'''} \cdot \nabla z_z''' - \overline{\omega}^e), \quad (10b)$$

where (4b) has been used.⁴ The expressions in brackets will prove useful in section 3. As noted in AG12, the quasi-Stokes velocity and the Stokes drift are closely related.

b. A new expression of the form stress term

The form stress term in (8) is based on the pressure fluctuation p''' so that it is not useful for some analytical treatments (Arduin et al. 2008a,b). Our view is that writing the form stress term in the TWM momentum

equation (6b) in terms of pressure [i.e., as in (8)] implies a wave-averaged equation, which corresponds to the Lagrangian average of (1): the direct expression for the momentum equation in the three-dimensional Lagrangian coordinates with the complicated pressure gradient term. We now show that the pressure-based form stress term in (8) can be transformed into a new expression where the pressure fluctuation p''' does not appear and which, in turn, provides a link to the Lagrangian average of the transformed expression for the Lagrangian momentum equation (2).

Substitution of $z^e = z + z'''$ and $\mathcal{D}_t \equiv \partial_t + \mathbf{V} \cdot \nabla + \omega \partial_z$ to (4a)–(4d) yields

$$\mathcal{D}_t \mathbf{V} = -\nabla(p + \eta) - \underbrace{(\mathcal{D}_t w)}_{-p_{z^e}} \nabla z''' \quad \text{and} \quad (11a)$$

$$\mathcal{D}_t w = -p_z - (\mathcal{D}_t w) z_z'''. \quad (11b)$$

We derived (11b) by multiplying (4d) with $1 + z_z'''$ and noting that $(1 + z_z''')p_{z^e} = z_z^e p_{z^e} = p_z$. Using (11a)–(11b), we rewrite the form stress term (8) to give

$$\begin{aligned}\mathcal{F}S^{\mathbf{V}} &= -\overline{z_z'''\nabla(p'''+\eta''')} + \overline{(\nabla z''')p_z'''} \\ &= -\overline{z_z'''\nabla(p+\eta)} + \overline{(\nabla z''')p_z} \\ &= \overline{z_z'''}(\mathcal{D}_t \mathbf{V} + (\mathcal{D}_t w)\nabla z''') - \overline{\nabla z'''(1+z_z''')(\mathcal{D}_t w)} \\ &= \overline{z_z'''}(\mathcal{D}_t \mathbf{V}) - \overline{\nabla z'''}(\mathcal{D}_t w) \\ &= \overline{z_z'''}(\mathcal{D}_t \mathbf{V})''' - \overline{\nabla z'''}(\mathcal{D}_t w)''',\end{aligned}\quad (12)$$

where no p''' appears at the last line. This expression has not, to our knowledge, been shown before the present study and is the cornerstone of the present study.⁵ Note that (12) has been derived from a nonlinear equation system and thus is applicable to finite-amplitude waves including the Doppler effect by mean flows.

c. Manipulation of the velocity-based form stress term

To expand the velocity-based form stress term (12), we need to derive variants of (4a)–(4b). Using $z_z^e = 1 + z_z'''$ and $A'' \equiv A - \hat{A}$ for an arbitrary quantity, (4a) may be written

$$\begin{aligned}\mathcal{D}_t z_z''' &= -z_z^e[\mathbf{V} \cdot (\hat{\mathbf{V}} + \mathbf{V}'') + (\hat{\omega} + \omega'')]_z \\ &= -z_z^e(\mathbf{V} \cdot \mathbf{V}'' + \omega_z''),\end{aligned}\quad (13)$$

where (6a) has been used to derive the second line. The material derivative operator may be expanded as

⁴ Appendix D of AG13 provides an approximate expression for EM quantities based on a Taylor expansion in the vertical direction with a continuous treatment of the vertical profile (i.e., no singular treatment of the vicinity of the sea surface), and is free from the traditional issue of how to handle regions above the surface troughs.

⁵ Apart from the simplified version shown by AG13 at their (26).

$$\begin{aligned}
 \mathcal{D}_t &\equiv \partial_t + \mathbf{V} \cdot \nabla + \varpi \partial_z \\
 &= \partial_t + (\hat{\mathbf{V}} + \mathbf{V}'') \cdot \nabla + (\hat{\omega} + \varpi'') \partial_z \\
 &= \hat{\mathcal{D}}_t + \mathbf{V}'' \cdot \nabla + \varpi'' \partial_z.
 \end{aligned} \tag{14}$$

Substitution of (14) to (13) yields

$$\hat{\mathcal{D}}_t z_z''' = -\nabla \cdot (z_z^e \mathbf{V}'') - (z_z^e \varpi'')_z. \tag{15}$$

On the other hand, (4b) may be written using (14) and $z^e = z + z''$, as

$$\begin{aligned}
 w &= (\partial_t + \mathbf{V} \cdot \nabla + \varpi \partial_z) z^e \\
 &= \hat{\mathcal{D}}_t z^e + (\mathbf{V}'' \cdot \nabla + \varpi'' \partial_z) z^e \\
 &= \hat{\mathcal{D}}_t z_z''' + \hat{\omega} + \mathbf{V}'' \cdot \nabla z_z''' + \varpi'' z_z^e.
 \end{aligned} \tag{16}$$

The unweighted average of (16) yields

$$\bar{w} = \hat{\omega} + \overline{\mathbf{V}'' \cdot \nabla z_z''}. \tag{17}$$

Combining (17) and (16) then yields

$$w''' = \hat{\mathcal{D}}_t z_z''' + (\mathbf{V}'' \cdot \nabla z_z''')''' + \varpi'' z_z^e. \tag{18}$$

The quantity ϖ'' is sufficiently small (as we shall see later), that all terms containing ϖ'' in the above are neglected in what follows.

We are now ready to expand the terms that make up the last line of (12). Using (14) and (15), the first term on the last line of (12), letting A be either u or v , can be written

$$\begin{aligned}
 \overline{z_z''' (\mathcal{D}_t A)'''} &= \overline{z_z''' (\hat{\mathcal{D}}_t A)'''} \\
 &= \overline{z_z''' (\hat{\mathcal{D}}_t A + \mathbf{V}'' \cdot \nabla A)} \\
 &= \hat{\mathcal{D}}_t \overline{(z_z''' A)} - \overline{(\hat{\mathcal{D}}_t z_z''') A} + \overline{z_z''' \mathbf{V}'' \cdot \nabla A} \\
 &= \hat{\mathcal{D}}_t \overline{(z_z''' A)} + \overline{[\nabla \cdot (z_z^e \mathbf{V}'')] A} + \overline{z_z''' \mathbf{V}'' \cdot \nabla A} \\
 &= \hat{\mathcal{D}}_t \overline{(z_z''' A)} + \nabla \cdot \overline{(z_z^e \mathbf{V}'' A)} - \overline{z_z^e \mathbf{V}'' \cdot \nabla A} \\
 &\quad + \overline{z_z''' \mathbf{V}'' \cdot \nabla A} \\
 &= \hat{\mathcal{D}}_t \overline{(z_z''' A)} + \nabla \cdot \overline{(z_z^e \mathbf{V}'' A)} - \overline{\mathbf{V}'' \cdot \nabla A} \\
 &= \hat{\mathcal{D}}_t \overline{(z_z''' A)} + \nabla \cdot \overline{(z_z^e \mathbf{V}'' A)} - \overline{(\mathbf{V}''' - z_z''' \mathbf{V}''')} \cdot \nabla A \\
 &= \hat{\mathcal{D}}_t \overline{(z_z''' A)} + \mathcal{R}S^A - \overline{\mathbf{V}''' \cdot \nabla A}''' + \overline{z_z''' \mathbf{V}'''} \cdot \nabla \bar{A},
 \end{aligned} \tag{19}$$

where $\mathbf{V}''' \equiv \mathbf{V} - \hat{\mathbf{V}} = (\bar{\mathbf{V}} + \mathbf{V}''') - \hat{\mathbf{V}} = \mathbf{V}''' - \overline{z_z''' \mathbf{V}'''}$ has been used to derive the second last line, and $\nabla \cdot [z_z^e \mathbf{V}'' (A + A'')] = \nabla \cdot (z_z^e \mathbf{V}'' A'') = \mathcal{R}S^A$ has been used to derive the last line.

Turning now to the second term on the last line of (12), we use (14) and (18), letting subscript X correspond to either ∂_x or ∂_y , to write

$$\begin{aligned}
 -\overline{z_X''' (\mathcal{D}_t w)'''} &= -\overline{z_X''' (\hat{\mathcal{D}}_t w)'''} \\
 &= -\overline{z_X''' (\hat{\mathcal{D}}_t w + \mathbf{V}'' \cdot \nabla w)} \\
 &= -\hat{\mathcal{D}}_t \overline{(z_X''' w)} + \overline{(\hat{\mathcal{D}}_t z_X''') w} - \overline{z_X''' \mathbf{V}'' \cdot \nabla w} \\
 &= -\hat{\mathcal{D}}_t \overline{(z_X''' w)} + \overline{w_X''' w} - \hat{\mathbf{V}}_X \cdot \overline{(\nabla z_X''') w} \\
 &\quad - \hat{\omega}_X \overline{z_X''' w} - \overline{(\mathbf{V}'' \cdot \nabla z_X''') w} - \overline{z_X''' \mathbf{V}'' \cdot \nabla w} \\
 &= -\hat{\mathcal{D}}_t \overline{(z_X''' w)} - \bar{\pi}_X - \hat{\mathbf{V}}_X \cdot \overline{(\nabla z_X''') w} \\
 &\quad - \hat{\omega}_X \overline{z_X''' w} + \overline{(\mathbf{V}'' \cdot \nabla z_X''') w_X'''} - \overline{z_X''' \mathbf{V}'' \cdot \nabla w} \\
 &= -\hat{\mathcal{D}}_t \overline{(z_X''' w_X''')} - \bar{\pi}_X - \hat{\mathbf{V}}_X \cdot \overline{(\nabla z_X''') w_X'''} \\
 &\quad - \hat{\omega}_X \overline{z_X''' w_X'''} - \overline{z_X''' \mathbf{V}'' \cdot \nabla w} \\
 &\quad + \overline{\mathbf{V}'' \cdot (w_X''' \nabla z_X''') - z_X''' \nabla w_X'''}),
 \end{aligned} \tag{20}$$

where $\bar{\pi} \equiv -\overline{w'' w_X''} / 2 + \overline{(\mathbf{V}'' \cdot \nabla z_X''') w''}$ is the Bernoulli head. We developed (20) from AM78 (appendix A). On the other hand, with the vertical component of the momentum equation in mind, we substitute $A = w$ to (19) and $X = z$ to (20), and take the sum of the two equations to give

$$\begin{aligned}
 0 &= \mathcal{R}S^w - \overline{\mathbf{V}''' \cdot \nabla w}''' + \overline{z_z''' (\mathbf{V}''' - \mathbf{V}''')} \cdot \nabla \bar{w} - \bar{\pi}_z \\
 &\quad - \hat{\mathbf{V}}_z \cdot \overline{(\nabla z_z''') w}''' - \hat{\omega}_z \overline{z_z''' w}''' \\
 &\quad + \overline{\mathbf{V}'' \cdot (w_z''' \nabla z_z''') - z_z''' \nabla w_z'''}).
 \end{aligned} \tag{21}$$

The overall transformation in this section has been done without restricting the characteristics of the waves, and thus allows for finite-amplitude inhomogeneous unsteady waves.

3. Deriving the pseudomomentum in the VL framework

In this section, we show that the velocity-based expression for the form stress derived in the previous section contains the time derivative of a wave-induced velocity, which might be called the pseudomomentum in the VL framework, as it is analogous to the generalized pseudomomentum in AM78. When the velocity-based expression of the form stress term is combined with the TWM momentum equations (6b) and (6c), we obtain an expression for the time development of “the total transport velocity minus the VL pseudomomentum,” which might be called the quasi-EM velocity in the VL framework.

Because our aim in this section is to show that our equation for the time development of the quasi-EM velocity in the VL framework includes the vortex force, we adopt the same scaling for the waves and the

TABLE 3. Comparison of the scalings of the low-pass-filtered equations where $\alpha \ll 1$ is the surface slope of waves.

Equation system	Section 3 (23a)–(23c)	Appendix C (C2a)–(C2c)	Appendix C (C4a)–(C4c)	Section 4 (C4a), (C4c), and (49)
Coefficient of ∂_T	α^2	α^{n+2}	α^n	α^4
Coefficient of $\bar{\mathbf{V}}$	—	α^n	α^n	α^4
$\hat{\mathbf{V}}, \bar{\mathbf{V}}, \bar{\mathbf{V}}^e$	α^2	α^2	α^2	α^2
$\hat{w}, \bar{w}, \bar{w}^e$	α^2	α^{n+2}	α^{n+2}	α^6
Horizontal momentum equation	α^4	α^{n+4}	α^{n+2}	α^6
Vertical momentum equation	α^4	α^4	α^2	α^2
$\nabla \bar{\eta}$	α^4	α^{n+4}	α^{n+2}	α^6
ϖ''	α^5	α^{n+5}	α^{n+3}	α^7
$f\sigma$	—	—	—	α^4
$\nu\kappa^2/\sigma$	—	—	—	α^4

low-pass-filtered flow (i.e., LCs) as in CL76. Let $\alpha \ll 1$ be a measure of the surface slope of waves. CL76 assumed that (i) $O(\alpha)$ waves are steady and monochromatic, (ii) the strength of the low-pass-filtered flow is $O(\alpha^2)$, (iii) there is no separation between the wavelength of the waves and the horizontal scale of variation of the low-pass-filtered flow, and that (iv) the time development of the low-pass-filtered flow is two orders, in terms of α , slower than the phase cycle of the waves. The time derivative operator may then be decomposed as $\partial_t = \partial_\tau + \alpha^2 \partial_T$ where ∂_τ and ∂_T operates on wave and low-pass-filtered quantities, respectively. These conditions are summarized in Table 3. Throughout the remainder of the manuscript we assume an infinitely deep ocean. [We have confirmed that the machinery of the present study is also applicable to a flat-bottomed ocean (not shown). See also footnote 6.]

a. Asymptotic expansion

To be consistent, the mean component of all quantities is scaled at $O(\alpha^2)$ except for \bar{p} and $\bar{\eta}$, which are scaled at $O(\alpha^4)$,

$$\bar{z}^e \equiv z, \quad (22a)$$

$$\bar{\eta} = \alpha^4 \bar{\eta}_4 + O(\alpha^5), \quad (22b)$$

$$\bar{p} = \alpha^4 \bar{p}_4 + O(\alpha^5), \quad (22c)$$

$$\bar{\mathbf{V}} = \alpha^2 \bar{\mathbf{V}}_2 + O(\alpha^3), \quad (22d)$$

$$\bar{w} = \alpha^2 \bar{w}_2 + O(\alpha^3), \quad \text{and} \quad (22e)$$

$$\hat{w} = \alpha^2 \hat{w}_2 + O(\alpha^3). \quad (22f)$$

The numeric subscripts represent the order of an asymptotic expansion, which is as in AG12 and AG13. Then we specialize to the $O(\alpha^2)$ terms in the TWM incompressibility (6a) as well as the $O(\alpha^4)$ terms in the TWM momentum equations (9a)–(9b) to yield

$$\nabla \cdot \hat{\mathbf{V}}_2 + \hat{w}_{2z} = 0, \quad (23a)$$

$$\hat{D}_T \hat{\mathbf{V}}_2 + \mathcal{R}S_4^{\mathbf{V}} = -\nabla(\bar{p}_4 + \bar{\eta}_4) + \mathcal{F}S_4^{\mathbf{V}}, \quad \text{and} \quad (23b)$$

$$\hat{D}_T \hat{w}_2 + \mathcal{R}S_4^w = -\bar{p}_{4z}, \quad (23c)$$

where $\hat{D}_T \equiv \partial_T + \hat{\mathbf{V}}_2 \cdot \nabla + \hat{w}_2 \partial_z$ is the material derivative operator based on the total transport velocity at $O(\alpha^2)$. On the other hand, the fluctuation component of all quantities is expanded from $O(\alpha)$, except for ϖ'' which is expanded from $O(\alpha^5)$

$$z''' = \alpha z_1''' + \alpha^2 z_2''' + \alpha^3 z_3''' + O(\alpha^4), \quad (24a)$$

$$\eta''' = \alpha \eta_1''' + \alpha^2 \eta_2''' + \alpha^3 \eta_3''' + O(\alpha^4), \quad (24b)$$

$$p''' = \alpha p_1''' + \alpha^2 p_2''' + \alpha^3 p_3''' + O(\alpha^4), \quad (24c)$$

$$\mathbf{V}''' = \alpha \mathbf{V}_1''' + \alpha^2 \mathbf{V}_2''' + \alpha^3 \mathbf{V}_3''' + O(\alpha^4), \quad (24d)$$

$$w''' = \alpha w_1''' + \alpha^2 w_2''' + \alpha^3 w_3''' + O(\alpha^4), \quad \text{and} \quad (24e)$$

$$\varpi'' = O(\alpha^5). \quad (24f)$$

The scaling of ϖ'' stems from the scaling of mean sea surface height $\bar{\eta}$. This is because the kinematic boundary condition at the sea surface is $\varpi'' = \mathbf{V}'' \cdot \nabla \bar{\eta}$ (AG12) so that to the leading order $\varpi_5'' = \mathbf{V}_1'' \cdot \nabla \bar{\eta}_4$ (or $\varpi_5'' = \mathbf{V}_1''' \cdot \nabla \bar{\eta}_4$ because $\mathbf{V}_1''' = \mathbf{V}_1''$).⁶ Therefore the asymptotic

⁶In contrast to the present study, the quantity $\varpi'' (=w^{*''})$ has been scaled at $O(\alpha^2)$ in AG13 because they considered waves in shallow water on a bottom slope of $O(\alpha)$. See (6a)–(6c) of AG13 for the explicit expressions of the kinematic boundary conditions at a sloping bottom in the VL framework (which should be compared with bottom boundary conditions in the three-dimensional Lagrangian framework in a future study). As shown by AG13, it is certainly possible to manipulate $\varpi'' (=w^{*''})$ even if it has been scaled at $O(\alpha^2)$. However, for simplicity, the present study assumes an infinitely deep ocean (or a flat-bottomed ocean) and thus neglects ϖ'' from (19) and thereafter.

TABLE 4. The rule of numeric subscript in the present study, which represents summation for a given order of asymptotic expansion in terms of α .

$(AB)_2 = A_1B_1$
$(AB)_3 = A_1B_2 + A_2B_1$
$(AB)_4 = A_1B_3 + A_2B_2 + A_3B_1$
$(ABC)_4 = A_2B_1C_1 + A_1B_2C_1 + A_1B_1C_2$

expansion for ϖ'' starts from $O(\alpha^5)$, allowing us to formally ignore the last term of (7) to give

$$\mathcal{RS}_4^{\mathbf{V}} = \nabla \cdot (\overline{\mathbf{V}''\mathbf{V}''})_4 + \nabla \cdot (\overline{z_z'''\mathbf{V}''\mathbf{V}''})_4 \quad \text{and} \quad (25a)$$

$$\mathcal{RS}_4^w = \nabla \cdot (\overline{\mathbf{V}''w''})_4 + \nabla \cdot (\overline{z_z'''\mathbf{V}''w''})_4. \quad (25b)$$

The numeric subscript attached to the brackets represents summation of terms at a given order of α (see Table 4 for a template). To obtain the $O(\alpha^4)$ form stress term for use in (23b), we use the velocity-based expression (12) to give

$$\mathcal{FS}_4^{\mathbf{V}} = \overline{[z_z'''\mathcal{D}_t\mathbf{V}'' - \nabla z_z'''\mathcal{D}_t w'']}_4. \quad (26)$$

To summarize, both the Reynolds stress term and the form stress term consist of the effect of waves up to $O(\alpha^3)$, as can be seen from Table 4.

Substitution of (22a)–(22f) and (24a)–(24f) to (4a)–(4d) yields

$$z_{1z\tau}''' + \nabla \cdot \mathbf{V}_1''' = 0, \quad (27a)$$

$$z_{1\tau}''' = w_1''', \quad (27b)$$

$$\mathbf{V}_{1\tau}''' = -\nabla(p_1''' + \eta_1'''), \quad \text{and} \quad (27c)$$

$$w_{1\tau}''' = -p_{1z}''', \quad (27d)$$

at $O(\alpha)$ and

$$z_{2z\tau}''' + \nabla \cdot (\mathbf{V}_2''' + z_{1z}'''\mathbf{V}_1''') = 0, \quad (28a)$$

$$z_{2\tau}''' + \mathbf{V}_1''' \cdot \nabla z_1''' = w_2''', \quad (28b)$$

$$\mathbf{V}_{2\tau}''' + \mathbf{V}_1''' \cdot \nabla \mathbf{V}_1''' = -\nabla(p_2''' + \eta_2''') - w_{1\tau}'''\nabla z_1''', \quad \text{and} \quad (28c)$$

$$w_{2\tau}''' + \mathbf{V}_1''' \cdot \nabla w_1''' = -p_{2z}''' - w_{1\tau}'''\nabla z_1''', \quad (28d)$$

at $O(\alpha^2)$. Note that because the first-order waves are steady and horizontally homogeneous, it follows from (28a)–(28d) that the second-order waves are also steady and horizontally homogeneous. The effect of the low-pass-filtered flow (i.e., LCs) on the waves appears in

equations for $O(\alpha^3)$ waves, to be explained later in the manuscript.

b. First- and second-order waves

As assumed above, $O(\alpha)$ waves are monochromatic and steady: $\eta_1''' = \mathcal{A} \cos\theta$ where \mathcal{A} is wave amplitude, $\theta = kx + ly - \sigma\tau$ is wave phase with k and l being wave-numbers in the x and y direction, and σ is wave frequency. These parameters are constant on the time and spatial scales of waves (i.e., $\partial_\tau \mathcal{A} = 0$ and $\nabla \mathcal{A} = 0$ for $\mathcal{A} = \mathcal{A}, k, l, \sigma$), which leads to

$$\partial_\tau (\overline{A_1'''B_1'''}) = 0, \quad \overline{A_1'''B_{1\tau}'''} = -\overline{A_{1\tau}'''B_1'''} \quad \text{and} \quad (29a)$$

$$\nabla (\overline{A_1'''B_1'''}) = 0, \quad \overline{A_1'''\nabla B_1'''} = -\overline{(\nabla A_1''')}B_1''', \quad (29b)$$

where A_1''' and B_1''' are wave variables at $O(\alpha)$. The sea surface is located at $z^\varepsilon = \bar{\eta} + \eta_1'''$ in the Eulerian–Cartesian coordinates and is located at $z = \bar{\eta} = \alpha^4 \bar{\eta}_4 + O(\alpha^5)$ in the VL coordinates. As far as up to $O(\alpha^3)$ waves are concerned, $z = 0$ can be used as the label of sea surface in the VL coordinates. With the boundary conditions of $\mathbf{V}_1''' = 0$ at $z = -\infty$ and $p_1''' = 0$ at $z = 0$, we solve (27a)–(27d) to yield

$$\sigma^2 = \kappa, \quad \kappa \equiv \sqrt{k^2 + l^2}, \quad (30a)$$

$$\phi_1''' \equiv (\mathcal{A}/\kappa)(\exp\kappa z) \cos\theta, \quad (30b)$$

$$\mathbf{V}_1''' = \nabla \phi_{1\tau}''' = \sigma(\nabla\theta)\phi_1''', \quad (30c)$$

$$w_1''' = \phi_{1z\tau}''' = -\sigma\phi_{1z\theta}''', \quad (30d)$$

$$z_1''' = \phi_{1z}''', \quad \text{and} \quad (30e)$$

$$p_1''' = \sigma^2\phi_1''' - \eta_1''', \quad (30f)$$

where $\nabla\phi_1''' = (\nabla\theta)\phi_{1\theta}''', \phi_{1\tau}''' = -\sigma\phi_{1\theta}''', \nabla\theta = (k, l)$, and $\eta_1''' = z_1'''|_{z=0}$ are understood. The above solution is given in the VL coordinates. Then, we compute the quasi-Stokes velocity using (10a)–(10b). For an arbitrary quantity A , the TWM and the EM at $O(\alpha^2)$ can be written as

$$\hat{A}_2 = \bar{A}_2 + \overline{z_{1z}'''\mathcal{A}_1'''} \quad \text{and} \quad (31a)$$

$$\bar{A}_2^\varepsilon = \bar{A}_2 - \overline{z_1'''\mathcal{A}_{1z}'''}, \quad (31b)$$

where (31b) is given by a Taylor expansion in the vertical direction (see footnote 4). Substitution of (30c)–(30e) and (31a)–(31b) to (10a)–(10b) yields

$$\mathbf{V}_2^{\text{qs}} = \overline{z_{1z}'''\mathbf{V}_1'''} + \overline{z_1'''\mathbf{V}_{1z}'''} = \overline{(z_1'''\mathbf{V}_1''')}_{z=0} = \overline{(\phi_{1z}'''\phi_1''')}_{z=0} \sigma \nabla\theta \quad \text{and} \quad (32a)$$

$$w_2^{\text{qs}} = -\overline{\mathbf{V}_1'' \cdot \nabla z_1'''} + \overline{z_1'''} w_{1z}'' = -\mathbf{V} \cdot (\overline{z_1'''} \mathbf{V}_1''') = 0, \quad (32b)$$

which has been shown by Mellor (2003) and Smith (2006), except that these authors did not refer to the term ‘‘quasi-Stokes velocity’’. In (32b), the condition of horizontally homogeneous waves leads to $w_2^{\text{qs}} = 0$, namely, $\hat{w}_2 = \overline{w_2^e}$.

Substitution of the $O(\alpha)$ solution (30c)–(30e) to $O(\alpha^2)$ momentum equations (28c)–(28d) yields

$$\mathbf{V}_{2\tau}''' = -\nabla(p_2''' + \eta_2''') \quad \text{and} \quad (33a)$$

$$w_{2\tau}''' = -\partial_z(p_2''' + \eta_2'''), \quad (33b)$$

which indicates that $O(\alpha^2)$ velocity [as well as $O(\alpha)$ velocity] satisfies an apparent irrotational condition in the VL coordinates:

$$\nabla \times \mathbf{V}_i''' = 0 \quad \text{and} \quad \nabla w_i''' - \partial_z \mathbf{V}_i''' = 0, \quad (34)$$

for $i = 1$ and 2 , where here, $\nabla \times \mathbf{V}''' = (v_x''' - u_y''')\mathbf{z}$, and \mathbf{z} is a unit vector in the upward vertical direction.⁷

c. Substitution of the velocity-based form stress term

The condition of horizontally homogeneous and steady waves is limited to up to $O(\alpha^2)$. The Doppler effect by both the horizontal and vertical circulations associated with the low-pass-filtered flow appears when considering the waves at $O(\alpha^3)$. However, it is rather difficult to derive (i) an analytical solution for $O(\alpha^3)$ waves and (ii) a depth-dependent wave crest equation. It is the depth-independent wave crest equation that has been used in Garrett (1976) and Smith (2006). Leibovich (1980) avoids these difficulties using the Lagrangian mean framework, an approach we mimic here. In fact the two terms on the last line of (12) have been expanded using (4a) and (4b), respectively, to give (19) and (20) (see section 2c). We pick up the $O(\alpha^4)$ terms of (19)–(20) and delete some terms, using the phase relationship of $O(\alpha)$ waves in (30c)–(30e), to give

$$\begin{aligned} \mathcal{FS}_4^u &= \hat{D}_T(\overline{z_{1z}'''} u_1'' - \overline{z_{1x}'''} w_1''') + \overline{z_{1z}'''} \mathbf{V}_1''' \cdot \nabla \bar{u}_2 \\ &\quad - \hat{\mathbf{V}}_{2x} \cdot (\overline{\nabla z_1'''} w_1'' + \mathcal{RS}_4^u - \bar{\pi}_{4x} - (\overline{\mathbf{V}''' \cdot \nabla u''}))_4 \end{aligned}$$

and

$$(35a)$$

$$\begin{aligned} \mathcal{FS}_4^v &= \hat{D}_T(\overline{z_{1z}'''} v_1'' - \overline{z_{1y}'''} w_1''') + \overline{z_{1z}'''} \mathbf{V}_1''' \cdot \nabla \bar{v}_2 \\ &\quad - \hat{\mathbf{V}}_{2y} \cdot (\overline{\nabla z_1'''} w_1'' + \mathcal{RS}_4^v - \bar{\pi}_{4y} - (\overline{\mathbf{V}''' \cdot \nabla v''}))_4, \end{aligned} \quad (35b)$$

where $\bar{\pi}_4 \equiv -(\overline{w''^2})_4/2 + [(\overline{\mathbf{V}'' \cdot \nabla z'''} w''')]_4$ is the Bernoulli head (to be updated later in the manuscript).⁸ Substitution of (35a)–(35b) to (23b) yields

$$\begin{aligned} \hat{D}_T(\bar{u}_2 + \overline{z_{1x}'''} w_1''') &= -(\bar{p}_4 + \bar{\eta}_4 + \bar{\pi}_4)_x + \overline{z_{1z}'''} \mathbf{V}_1''' \cdot \nabla \bar{u}_2 \\ &\quad - \hat{\mathbf{V}}_{2x} \cdot (\overline{\nabla z_1'''} w_1'' - (\overline{\mathbf{V}''' \cdot \nabla u''}))_4 \quad \text{and} \end{aligned} \quad (36a)$$

$$\begin{aligned} \hat{D}_T(\bar{v}_2 + \overline{z_{1y}'''} w_1''') &= -(\bar{p}_4 + \bar{\eta}_4 + \bar{\pi}_4)_y + \overline{z_{1z}'''} \mathbf{V}_1''' \cdot \nabla \bar{v}_2 \\ &\quad - \hat{\mathbf{V}}_{2y} \cdot (\overline{\nabla z_1'''} w_1'' - (\overline{\mathbf{V}''' \cdot \nabla v''}))_4, \end{aligned} \quad (36b)$$

where the effect of $O(\alpha^3)$ waves appear in both the Bernoulli head $\bar{\pi}_4$ and the last term of each equation. The last term of (36a)–(36b) is the legacy of the Reynolds stress term and is absent in the three-dimensional Lagrangian framework of AM78. It should be noted that the $\hat{D}_T \hat{\mathbf{V}}_2$ term on the lhs of (23b) has been partially cancelled by the \hat{D}_T terms on the rhs of (35a)–(35b) using $\hat{\mathbf{V}}_2 = \nabla_2 + \overline{z_{1z}'''} \mathbf{V}_1'''$, which follows from (31a). This is the first indication of the appearance of the pseudomomentum, to be discussed in detail in the next subsection.

A nice feature of AM78 is that the Bernoulli head is present in both the horizontal and vertical components of the wave-averaged momentum equations, and there is no need to treat the Bernoulli head and nonhydrostatic pressure separately (cf. Craik 1985; Dingemans 2009). The sum of the Bernoulli head and nonhydrostatic pressure can be obtained by solving a Poisson equation based on the incompressibility condition of circulation (as is always the case in nonhydrostatic numerical models). Eventually there is no need to derive the analytical solution for $O(\alpha^3)$ waves. It is therefore very useful to mimic this feature of AM78—the Bernoulli head appearing next to nonhydrostatic pressure in both the horizontal and vertical components of the wave-averaged momentum equations. So far the Bernoulli head in our analysis appears only in the horizontal component of the TWM momentum equations (36a)–(36b). However, it is

⁷ Although (34) is sufficient for us to derive the vortex force, the standard vorticity defined in the three-dimensional Eulerian coordinates reads $\nabla^e \times \mathbf{V} = (z_z^e \nabla \times \mathbf{V} - \nabla z^e \times \mathbf{V}_z)/z_z^e$ and $\nabla^e w - \mathbf{V}_{ze} = (z_z^e \nabla w - w_z \nabla z^e - \mathbf{V}_z)/z_z^e$.

⁸ The last term on the last line of (20) becomes $\overline{v''(\nabla w'' \times \nabla z''')}$ in the case of $X = x$, and $-\overline{u''(\nabla w'' \times \nabla z''')}$ in the case of $X = y$. When taking an asymptotic expansion of these terms, $\nabla w_i''' \times \nabla z_j''' = 0$ for $(i, j) = (1, 1)$, $(1, 2)$, and $(2, 1)$. This is because $\nabla w_i''' \propto \nabla \theta$ and $\nabla z_i''' \propto \nabla \theta$ according to the analytical solution of $O(\alpha)$ and $O(\alpha^2)$ waves.

straightforward to write the vertical momentum equation in a form that includes the Bernoulli head. Indeed, we have derived (21), which allows us to transform the Reynolds stress term in the vertical component of the momentum equation (23c), into the sum of the vertical gradient of the Bernoulli head and the other terms, as follows:

$$\begin{aligned} \hat{D}_T \underbrace{(\bar{w}_2 + \overline{z_{1z}''' w_1'''})}_{\hat{w}_2} &= -(\bar{p}_4 + \bar{\pi}_4)_z - \hat{\mathbf{V}}_{2z} \cdot \overline{(\nabla z_1''') w_1'''} \\ &+ \overline{[\mathbf{V}'' \cdot (w_z''' \nabla z''' - z_z''' \nabla w''')]_4} \\ &- \overline{(\mathbf{V}''' \cdot \nabla w''')_4}, \end{aligned} \quad (36c)$$

$$\begin{pmatrix} \bar{u}_2 + \overline{z_{1x}''' w_1'''} \\ \bar{v}_2 + \overline{z_{1y}''' w_1'''} \\ \bar{w}_2 + \overline{z_{1z}''' w_1'''} \end{pmatrix} = \underbrace{\begin{pmatrix} \hat{u}_2 \\ \hat{v}_2 \\ \hat{w}_2 \end{pmatrix}}_{\text{total transport velocity}} - \underbrace{\begin{pmatrix} +z_{1z}''' & 0 & -z_{1x}''' \\ 0 & +z_{1z}''' & -z_{1y}''' \\ -z_{1x}''' & -z_{1y}''' & -z_{1z}''' \end{pmatrix} \begin{pmatrix} u_1''' \\ v_1''' \\ w_1''' \end{pmatrix}}_{\text{pseudomomentum}}, \quad (37)$$

where (31a) and (17) have been used. The wave-induced velocity on the rhs may be called the pseudomomentum in the VL framework, as it is analogous to the generalized pseudomomentum in AM78 (appendix A). By substituting (30c)–(30e) to the last term of (37), we confirm that the content of the pseudomomentum is identical to the quasi-Stokes velocity in (32a)–(32b). This association of the pseudomomentum in (37) to the quasi-Stokes velocity in (32a)–(32b) stems from the conditions of (i) apparent irrotational wave motions as given by (34) and (ii) horizontally homogeneous and steady waves as given by (29a)–(29b). In fact

$$\begin{aligned} \mathbf{V}_2^{\text{qs}} &= \overline{z_{1z}''' \nabla_1'''} + \overline{z_1''' \nabla_{1z}'''} \\ &= \overline{z_{1z}''' \nabla_1'''} + \overline{z_1''' \nabla w_1'''} \\ &= \overline{z_{1z}''' \nabla_1'''} - \overline{(\nabla z_1''') w_1'''} \quad \text{and} \end{aligned} \quad (38a)$$

$$\begin{aligned} w_2^{\text{qs}} &= -\overline{\mathbf{V}_1'' \cdot \nabla z_1'''} + \overline{z_1''' w_{1z}'''} \\ &= -\overline{\mathbf{V}_1'' \cdot \nabla z_1'''} + \overline{z_1''' z_{1z\tau}'''} \\ &= -\overline{\mathbf{V}_1'' \cdot \nabla z_1'''} - \overline{w_1''' z_{1z}'''}, \end{aligned} \quad (38b)$$

where (27b) has been used in (38b). The last line of each of (38a)–(38b) is identical to the definition of the pseudomomentum in (37). Recall that the difference of the total transport velocity and the quasi-Stokes velocity is the EM velocity. Hence the lhs of (36a)–(36c) is the same as $\hat{D}_T \bar{u}_2^e$, $\hat{D}_T \bar{v}_2^e$, and $\hat{D}_T \bar{w}_2^e$: the material derivative of the EM velocity.

where the expression for \hat{w}_2 uses (31a). Note that $O(\alpha^3)$ waves appear in both the Bernoulli head $\bar{\pi}_4$ and the last term. The second last term of (36c) consists of waves up to $O(\alpha^2)$ that are horizontally homogeneous (so that, as we shall show later, the term can be absorbed into the Bernoulli head without affecting the horizontal component of the momentum equations).

d. The pseudomomentum in the VL framework

Interestingly, the three-dimensional components of velocity on the lhs of (36a)–(36c) have a symmetry and can be written as the difference of the total transport velocity and a wave-induced velocity

To summarize, in this section we have shown how to extract the pseudomomentum in the VL framework from the expression for the form stress written in terms of velocity rather than pressure. It is now straightforward, although mathematically laborious, to derive the expression for the vortex force in the VL framework. This is done in appendix B.

4. The effect of viscosity

So far the analysis in the present study has been done for waves in a nonrotating inviscid fluid. As shown below, the velocity-based expression of the form stress term (12) is modified by the introduction of the Coriolis term and the viscosity term.

a. The form stress term for a rotating viscid fluid

Inclusion of the new terms to the momentum equations (11a)–(11b) in the VL coordinates yields

$$\mathcal{D}_t \mathbf{V} + f \mathbf{z} \times \mathbf{V} = -\nabla(p + \eta) - \underbrace{(\mathcal{D}_t w - F^w)}_{-p_{z^e}} \nabla z''' + F^{\mathbf{V}} \quad (39a)$$

and

$$\underbrace{(1 + z_z''')}_{z_z^e} \mathcal{D}_t w = -p_z + \underbrace{(1 + z_z''')}_{z_z^e} F^w, \quad (39b)$$

where $(1 + z_z''') p_{z^e} = z_z^e p_{z^e} = p_z$ is understood. The symbols $F^{\mathbf{V}}$ and F^w represent the effect of turbulent mixing

on \mathbf{V} and w , respectively. These terms are parameterized using a conventional symmetric tensor in Eulerian–Cartesian coordinates, as in (28) of AG12. The Coriolis parameter f as well as the coefficient of turbulent viscosity ν (to appear later) have been nondimensionalized following the approach of AG13.⁹

The TWM equation system (6a)–(6c) becomes

$$\mathbf{V} \cdot \hat{\nabla} + \hat{\omega}_z = 0, \quad (40a)$$

$$\begin{aligned} \hat{\mathbf{V}}_t + \nabla \cdot (\hat{\mathbf{V}}\hat{\mathbf{V}}) + (\hat{\omega}\hat{\mathbf{V}})_z + f\mathbf{z} \times \hat{\mathbf{V}} + \mathcal{R}S^{\mathbf{V}} \\ = -\nabla(\bar{p} + \bar{\eta}) + \mathcal{F}S^{\mathbf{V}} + \hat{F}^{\mathbf{V}}, \quad \text{and} \end{aligned} \quad (40b)$$

$$\hat{w}_t + \nabla \cdot (\hat{\mathbf{V}}\hat{w}) + (\hat{\omega}\hat{w})_z + \mathcal{R}S^w = -\bar{p}_z + \hat{F}^w, \quad (40c)$$

where the Coriolis term includes $f\mathbf{z} \times \mathbf{V}^{\text{qs}}$, which corresponds to the Coriolis–Stokes force of Hasselmann (1970) and Huang (1979). The Reynolds and form stress terms in (40b)–(40c) are the same as (7) and (8), respectively. The velocity-based expression of the form stress term (12) is revised as follows:

$$\begin{aligned} \mathcal{F}S^{\mathbf{V}} &= -\overline{z_z'''\nabla(p + \eta)} + \overline{(\nabla z''')p_z} \\ &= \overline{z_z'''\mathcal{D}_t\mathbf{V} + f\mathbf{z} \times \mathbf{V} + (\mathcal{D}_t w - F^w)\nabla z''' - F^{\mathbf{V}}} \\ &\quad - \overline{\nabla z'''(1 + z_z''')(\mathcal{D}_t w - F^w)} \\ &= \overline{z_z'''\mathcal{D}_t\mathbf{V}'''} - \overline{\nabla z'''(\mathcal{D}_t w)'''} + f\mathbf{z} \times \overline{z_z'''\mathbf{V}'''} \\ &\quad - \overline{z_z'''(F^{\mathbf{V}})'''} + \overline{(\nabla z''')(F^w)'''}, \end{aligned} \quad (41)$$

where (39a)–(39b) have been used.

b. Asymptotic expansion

The derivation leading to (41) was obtained without approximation. We now specialize to small-amplitude waves.

Jenkins (1987) investigated the problem of how the presence of surface waves modifies the classical Ekman spiral solution, and obtained an interesting result when the viscosity coefficient varies in the vertical direction (to be explained later). We wish to make the link to the work of Jenkins (1987), and thus retain the last three terms of (41) as follows. Some scalings typical for the ocean are $f/\sigma \sim O(\alpha^4)$ and $\nu\kappa^2/\sigma \sim O(\alpha^4)$ (see Table 2 of AG12 for the dimensional values of f , σ , κ , and ν). Thus the Coriolis parameter and the viscosity coefficient may

be scaled as $f = \alpha^4 f_4$ and $\nu = \alpha^4 \nu_4$, which indicates that the form stress term (41) should be written at $O(\alpha^6)$. For simplicity, we consider circulation whose variation is scaled as $m = n$ in appendix C and is associated with the equation system (C4a)–(C4c). Noting that the form stress term in (C4b) is written at $O(\alpha^{n+2})$, we obtain $m = n = 4$, namely $\partial_t = \partial_\tau + \alpha^4 \partial_T$ and $\nabla = \dot{\nabla} + \alpha^4 \bar{\nabla}$, where $\dot{\nabla}$ and $\bar{\nabla}$ are the lateral gradient operator for wave and low-pass-filtered quantities, respectively. See appendix C for details. These conditions are summarized in Table 3.

In what follows, we consider depths below the base of the thin viscous surface boundary layer (of a few centimeters depth) associated with waves (hereafter TVSBL). Thus, the $O(\alpha)$ quantities are the solution of inviscid waves (30a)–(30f). As in section 3, $O(\alpha)$ waves are assumed to be monochromatic, except that slow variations in both the horizontal direction and in time are allowed as in (C1a)–(C1b) (see footnote C1). We first note that the viscosity term of the TWM momentum equation (40b) is written by $\hat{F}_6^{\mathbf{V}} = \partial_z(\nu_4 \bar{\nabla}_{2z}^2)$, indicating that the viscosity acts on the EM velocity rather than the TWM velocity. This has been shown by (30) and (44) of AG12 and note that the result holds even if the viscosity coefficient varies in the vertical.

Coming back to the velocity-based form stress term, we substitute (19)–(20) to (41), and then pick-up $O(\alpha^6)$ terms to yield

$$\begin{aligned} \mathcal{F}S_6^{\mathbf{V}} &= \partial_T \mathbf{V}_2^{\text{qs}} + \mathcal{R}S_6^{\mathbf{V}} \\ &\quad - \frac{1}{2} \overline{\nabla(|\mathbf{V}_1'''|^2 - w_1'''^2)} - \overline{[(\nabla \times \mathbf{V}''') \times \mathbf{V}''']}_6 \\ &\quad + f_4 \mathbf{z} \times \overline{z_{1z}''' \mathbf{V}_1'''} - \overline{z_{1z}''' (F^{\mathbf{V}})_5'''} + \overline{(\dot{\nabla} z_{1z}''') (F^w)_5'''}, \end{aligned} \quad (42)$$

where (38a) and (B5a) have been used. The symbols $(F^{\mathbf{V}})_5'''$ and $(F^w)_5'''$ are the viscosity terms that are written using the solution of $O(\alpha)$ waves

$$\begin{aligned} (F^u)_5''' &= (2\nu_4 u_{1x}''')_x + [\nu_4 (v_{1x}''' + u_{1y}''')]_y + [\nu_4 (w_{1x}''' + u_{1z}''')]_z \\ &= \kappa\sigma [2\nu_4 \dot{\nabla}^2 \phi_1''' + (2\nu_4 \phi_{1z}''')]_z, \end{aligned} \quad (43a)$$

$$\begin{aligned} (F^v)_5''' &= [\nu_4 (u_{1y}''' + v_{1x}''')]_x + (2\nu_4 v_{1y}''')_y + [\nu_4 (w_{1y}''' + v_{1z}''')]_z \\ &= \iota\sigma [2\nu_4 \dot{\nabla}^2 \phi_1''' + (2\nu_4 \phi_{1z}''')]_z, \quad \text{and} \end{aligned} \quad (43b)$$

$$\begin{aligned} (F^w)_5''' &= [\nu_4 (u_{1z}''' + w_{1x}''')]_x + [\nu_4 (v_{1z}''' + w_{1y}''')]_y + (2\nu_4 w_{1z}''')_z \\ &= [2\nu_4 \dot{\nabla}^2 w_1''' + (2\nu_4 w_{1z}''')]_z, \end{aligned} \quad (43c)$$

⁹The nondimensionalization is written by $\hat{f} = (\hat{g}/\hat{\Delta})^{1/2} f$ and $\hat{\nu} = (\hat{\Delta}^3 \hat{g})^{1/2} \nu$ using the notation in appendix A of AG13.

where the second line of each has been derived using (30b)–(30d) and $\partial_x \nu_4 = \partial_y \nu_4 = 0$. The viscosity coefficient ν_4 is allowed to vary in the vertical direction, which is similar to Jenkins (1987).

We now consider how to calculate the fourth term on the rhs of (42), which consists of waves up to $O(\alpha^5)$. The solution of higher-order waves may be decomposed into that associated with the nonlinear terms of (39a)–(39b) and that associated with the effect of the Coriolis and viscosity terms. The former solution is written in terms of the harmonics of $O(\alpha)$ waves (not shown) and averages to zero and thus is not discussed further (this approach follows section 4b of AG13). Hereafter we focus on the latter solution, which is derived from the linear terms of (39a) to read

$$\partial_\tau \mathbf{V}_5''' + \partial_T \mathbf{V}_1''' + f_4 \mathbf{z} \times \mathbf{V}_1''' = -\dot{\mathbf{V}} p_5''' - \bar{\nabla} p_1''' + (F^{\mathbf{V}})_5''' . \quad (44)$$

We take the curl of (44) to yield,

$$\begin{aligned} \dot{\mathbf{V}} \times (\partial_\tau \mathbf{V}_5''') + \dot{\mathbf{V}} \times (f_4 \mathbf{z} \times \mathbf{V}_1''') &= -\dot{\mathbf{V}} \times \bar{\nabla} (p_1''' + \eta_1''') \\ &= \bar{\nabla} \times \dot{\mathbf{V}} (p_1''' + \eta_1''') \\ &= -\bar{\nabla} \times (\partial_\tau \mathbf{V}_1''') , \end{aligned} \quad (45)$$

where the last term of (44) has canceled out because $(F^{\mathbf{V}})_5''' \propto \dot{\mathbf{V}}\theta = (k, l)$, which follows from (43a)–(43b). The last line of (45) has been derived using (27c). Using (45), $\dot{\mathbf{V}} \cdot \mathbf{V}_1''' + w_{1z}''' = 0$ and $w_1''' = z_{1\tau}'''$, we show

$$\begin{aligned} [(\nabla \times \mathbf{V}''') \times \mathbf{V}''']_6 &= (\dot{\mathbf{V}} \times \mathbf{V}_5''') \times \mathbf{V}_1''' + (\bar{\nabla} \times \mathbf{V}_1''') \times \mathbf{V}_1''' \\ &= \int_\tau^\tau [\dot{\mathbf{V}} \times (\partial_\tau \mathbf{V}_5''') + \bar{\nabla} \times (\partial_\tau \mathbf{V}_1''')] d\tau \times \mathbf{V}_1''' \\ &= -\int_\tau^\tau \dot{\mathbf{V}} \times (f_4 \mathbf{z} \times \mathbf{V}_1''') d\tau \times \mathbf{V}_1''' \\ &= \int_\tau^\tau f_4 \mathbf{z} w_{1z}''' d\tau \times \mathbf{V}_1''' \\ &= f_4 \mathbf{z} \times z_{1z}''' \mathbf{V}_1''' , \end{aligned} \quad (46)$$

where the first line has been derived using $\dot{\mathbf{V}} \times \mathbf{V}_i''' = 0$ for $i = 1, 2, 3, 4$. Equation (46) indicates that the fourth and fifth terms on the rhs of (42) cancel each other.

We then calculate the viscosity terms of (42). The last two terms of (42) may be rewritten using $z_{1z}''' = \phi_{1zz}''' = \kappa^2 \phi_1''' = (\dot{\mathbf{V}}\theta \cdot \dot{\mathbf{V}}\theta) \phi_1''' = (1/\sigma)(\mathbf{V}_1''' \cdot \dot{\mathbf{V}}\theta)$ and $\dot{\mathbf{V}} z_{1\tau}''' = (\dot{\mathbf{V}}\theta) z_{1\tau}''' = (-1/\sigma)(\dot{\mathbf{V}}\theta) w_1'''$ to yield

$$\begin{aligned} & -z_{1z}''' (F^{\mathbf{V}})_5''' + \overline{(\dot{\mathbf{V}} z_{1\tau}''') (F^w)_5'''} \\ &= -\frac{1}{\sigma} \overline{(\mathbf{V}_1''' \cdot \dot{\mathbf{V}}\theta) (F^{\mathbf{V}})_5'''} - \frac{\dot{\mathbf{V}}\theta}{\sigma} \overline{w_1''' (F^w)_5'''} \\ &= -\frac{\dot{\mathbf{V}}\theta}{\sigma} \underbrace{\overline{[\mathbf{V}_1''' \cdot (F^{\mathbf{V}})_5'''] + w_1''' (F^w)_5''']}}_{\text{FluxDiv-Dissipation}} , \end{aligned} \quad (47)$$

where the last line has been derived using $(F^{\mathbf{V}})_5''' = (2\sigma \dot{\mathbf{V}}\theta)[\nu_4 \dot{\mathbf{V}}^2 \phi_1''' + (\nu_4 \phi_{1zz}''')]_z$, which follows from (43a)–(43b). Substitution of (46)–(47) to (42) yields

$$\begin{aligned} \mathcal{F}S_6^{\mathbf{V}} &= \partial_T \mathbf{V}_2^{\text{qs}} + \mathcal{R}S_6^{\mathbf{V}} - \frac{1}{2} \bar{\nabla} \cdot \overline{(|\mathbf{V}_1'''|^2 - w_1'''^2)} \\ & \quad - \frac{\dot{\mathbf{V}}\theta}{\sigma} \underbrace{\overline{[\mathbf{V}_1''' \cdot (F^{\mathbf{V}})_5'''] + w_1''' (F^w)_5''']}}_{\text{FluxDiv-Dissipation}} , \end{aligned} \quad (48)$$

where the last term is the product between $-(\dot{\mathbf{V}}\theta)/\sigma$ and the viscosity term of the depth-dependent energy equation, the latter of which may be separated into two terms: one is the vertical divergence of a viscosity-induced flux (noted as FluxDiv) and one is dissipation at depths below $z = \bar{\eta} - \delta$ (noted as Dissipation), where $\delta (>0)$ is the thickness of the TVSBL. This separation, in particular the identification of the dissipation term in the wave energy equation, is based on Phillips (1977, p. 52). See appendix D of the present manuscript for details. Substitution of (48) to either (40b) or the rotating viscid version of (C4b) yields

$$\partial_T \bar{\mathbf{V}}_2^e + f_4 \mathbf{z} \times \dot{\mathbf{V}}_2 = -\bar{\nabla} [\bar{p}_2 + \bar{\eta}_2 + (|\mathbf{V}_1'''|^2 - w_1'''^2)/2] + \underbrace{\partial_z (\nu_4 \bar{\mathbf{V}}_{2z}^e)}_{\hat{F}_6^{\mathbf{V}}} + \underbrace{\partial_z (-\nu_4 \mathbf{V}_{2z}^{\text{qs}})}_{(-\dot{\mathbf{V}}\theta/\sigma)\text{FluxDiv}} + \underbrace{\nu_4 \mathbf{V}_{2z}^{\text{qs}}}_{(\dot{\mathbf{V}}\theta/\sigma)\text{Dissipation}} , \quad (49)$$

where the last two terms have been derived using (32a) and (D1a)–(D1b).

The last term in (49) originates from the dissipation term in the wave energy equation (D2) and acts like a depth-dependent body force, indicating that the dissipation

of wave kinetic energy leads to transfer of momentum from waves to circulation (Fig. 2). The second to last term of (49) is associated with the vertical flux of kinetic energy in the depth-dependent wave energy equation. It is important to remember from the outset that the last




FIG. 2. Schematic of momentum transfer between wind, surface waves, and ocean circulation.

two terms on the rhs of (49) originate from the form stress term. As such, they represent the effect of the waves on circulation and have a vertical structure. In contrast to the present study, the explanation in Smith (2006) and Weber et al. (2006) is based on depth-integrated equations.

Once the last two terms of (49) are merged, the equation is identical to (5.1) of Jenkins (1987) who obtained it from the three-dimensional Lagrangian framework of Pierson (1962). Indeed, when the viscosity varies in the vertical, merging the last two terms of (49) leads to the “additional source of momentum” in the water column noted by Jenkins (1987). Nevertheless, the surface boundary condition derived by Jenkins (1987) is different from that appropriate to our study (see section 4c below). As explained in footnote B1 of AG12, we believe the boundary condition used by Jenkins (1987) [and also by Weber (1983)] is not correct.

c. The virtual wave stress of Longuet-Higgins (1953, 1960)

We now consider the surface boundary condition of the momentum equation (49), assuming that the net momentum flux through the TVSBL is vertically uniform. In doing so, we implicitly make use of the fact that the TVSBL, being only centimeters thick, is much thinner than the Ekman layer associated with the rotation of the earth, typically measurable in meters, or even tens of meters—see Table 2 in AG12.

Combining the vertical flux of momentum associated with the second and third last terms of (49) yields $\nu_4(\bar{\mathbf{V}}_2^e - \mathbf{V}_2^{qs})_z$ at the base of the TVSBL ($z = \bar{\eta} - \delta$). This momentum flux should match the skin viscous stress applied at the top of the TVSBL, namely the ocean

surface ($z = \bar{\eta}$).¹⁰ If there is no wind, the skin stress at the ocean surface is zero. Therefore the viscosity-induced momentum flux at the base of the TVSBL is also zero: $\nu_4(\bar{\mathbf{V}}_2^e - \mathbf{V}_2^{qs})_z = 0$ at $z = \bar{\eta} - \delta$. The result that $\bar{\mathbf{V}}_{2z}^e = \mathbf{V}_{2z}^{qs}$ is consistent with Longuet-Higgins (1953, 1960) who found that the vertical gradient of the LM velocity at the base of the TVSBL is twice that of the Stokes-drift velocity. It follows that the viscosity-induced stress $\nu_4 \mathbf{V}_{2z}^{qs}$ corresponds to the VWS. The explanation in the present study makes it clear that this result of Longuet-Higgins is more general than it might appear. In particular, the role played by the VWS has emerged without the explicit use of the analytical solution of waves in the TVSBL and thus is easily applicable to various types of problems. It does not matter whether there is wind forcing or not, the waves can have slow variations, and the viscosity coefficient can vary in the vertical—the result is quite general. Indeed, (49) represents an extension beyond the approach of Ünlüata and Mei (1970), Weber (1983), Xu and Bowen (1994), Ng (2004), and AG12 based on the analytical solution of waves including the TVSBL. The above explanation of the VWS also works for the scaling for the vortex force equations in section 3 (not shown).

Furthermore, (49) provides a prescription for including surface wave effects in ocean circulation models. At the sea surface ($z = \bar{\eta}$), the net momentum flux from air (i.e., wind) to water (i.e., ocean circulation and waves) is given by the sum of the skin stress and the wave stress (Fig. 2), the latter of which represents the residual effect of both the normal stress and the tangential stress associated with the waves (not shown, cf. Fan et al. 2010; Donelan et al. 2012; appendix B of AG12). The former (the skin stress) represents the direct transfer of momentum from wind to ocean circulation, and should match $\nu_4(\bar{\mathbf{V}}_2^e - \mathbf{V}_2^{qs})_z$ at the base of the TVSBL ($z = \bar{\eta} - \delta$). The latter (the wave stress) represents the transfer of momentum from wind to waves. The momentum of waves is eventually transferred to circulation when/where waves are dissipated by the turbulent viscosity, as is shown by the last term of (49). To summarize $\nu_4(\bar{\mathbf{V}}_2^e - \mathbf{V}_2^{qs})_z$ as “the skin stress” is the surface boundary condition applicable to numerical circulation models. The net momentum input to ocean circulation is given by the sum of the skin stress and the vertical integral of the last term of (49) associated with

¹⁰ The skin stress at the sea surface corresponds to the mean tangential stress $\bar{\tau}$ in AG12 (noting that τ in the present study has a different meaning and represents the time measure associated with the phase cycle of the waves). The definitions of the skin stress and the wave stress are partly related and require care, a topic we shall discuss in a later paper.

the wave dissipation. The net momentum input to traditional ocean circulation models (that is models that do not include wave effects) is given by the wind stress based, for example, on the Large and Pond (1981) parameterization, and should be compared with the sum of the skin stress and the depth integral of the wave dissipation term mentioned above.

5. Summary and discussion

The fundamentals of the vertically Lagrangian and horizontally Eulerian (VL) framework have been developed in the present study concerning the effect of surface waves on circulations in the upper ocean. We suggest that the thickness-weighted-mean (TWM) momentum equations of Mellor (2003), Broström et al. (2008), and AG12 correspond to the Lagrangian average of (1), which is the direct expression of the Lagrangian momentum equations. To our knowledge, no previous study has shown (i) how to derive the transformed expression for the TWM momentum equations corresponding to the Lagrangian average of (2) and (ii) how to introduce the concept of pseudomomentum to the VL framework, as can be seen in, for example, the discussion between Ardhuin et al. (2008a) and Mellor (2008b).

In section 2, we have shown that the traditional pressure-based form stress term can be transformed into a set of terms that do not contain any pressure quantities. The transformation in the present study is applicable to a nonlinear equation system, which is an improvement over AG13 who utilized a version of the transformation based on a linear equation system for the waves. An important by-product is that the velocity-based form stress term includes the time derivative of a wave-induced velocity which is referred to as the pseudomomentum in the VL framework, as it is analogous to the generalized pseudomomentum in AM78. The result is that the transformed expression of the TWM momentum equations (i.e., for the development of the quasi-EM velocity, namely the total transport velocity minus the VL pseudomomentum vector) has been obtained in section 3. As shown in appendix B, it is possible to derive the vortex force using the VL framework, using an approach that is a hybrid of Leibovich (1980) and CL76. We also noted that the twin expressions for the Lagrangian mean momentum equations (and hence also for the TWM momentum equations) may be traced back to the work of Lagrange (1788), which has been little mentioned in previous studies.

A nice feature of the VL framework is the treatment of the turbulent viscosity term near the sea surface. The traditional explanation for the viscosity-induced transfer

of momentum from waves to circulation has been based on depth-integrated equations (e.g., Smith 2006; Weber et al. 2006; Fan et al. 2010), whereas our explanation in section 4 is based on depth-dependent equations with a vertically nonuniform viscosity coefficient, and thus is useful for revisiting the surface boundary condition used in numerical circulation models. We have shown that the velocity-based expression of the form stress term contains the residual effect of viscosity [see (48)]. In the transformed expression of the TWM momentum equations, the effect of viscosity appears as the sum of a flux-divergence term (which is associated with the skin stress applied by wind) and a body force (which represents transfer of momentum from waves to circulation associated with the dissipation of wave kinetic energy) [see (49)]. This allows us to explain the concept of the virtual wave stress (VWS) of Longuet-Higgins (1953, 1960), without relying on the explicit use of the analytical solution of waves in the thin viscous boundary layer at the sea surface as in the work of Ünlüata and Mei (1970), Xu and Bowen (1994), Ng (2004), and AG12. Our explanation may be regarded as a recipe for a future study to explain the VWS using the three-dimensional Lagrangian framework of AM78, which has not been achieved in previous studies despite the utility of AM78 to allow a general spectrum of waves (cf. Ardhuin et al. 2008b).

Acknowledgments. HA thanks Bach Lien Hua for translating Lagrange (1788) and Hitoshi Tamura and Tetsu Hara for helpful discussions. RJG thanks GEOMAR for on-going support. We are also grateful to the anonymous reviewers of both the present and previous versions of this paper for their helpful comments.

APPENDIX A

Analogy to Andrews and McIntyre (1978)

The vector (Ξ_1, Ξ_2, Ξ_3) in AM78 represents the position of a fluid particle in the Eulerian–Cartesian coordinates, corresponding to (x^e, y^e, z^e) in the present study.^{A1} Likewise (x_1, x_2, x_3) in AM78 represents the position of a fluid particle in the three-dimensional Lagrangian coordinates, corresponding to (x, y, z) in the present study.^{A2} Thus the fluctuation of the position

^{A1} Numeric subscripts in AM78 represent axes in Cartesian coordinates and should not be confused with numeric subscripts in the present study representing the order of an asymptotic expansion.

^{A2} This has been written (a, b, c) in section 1 of the present study. In AM78, the coordinate transformation matrix has been written as $\Xi_{j,i} = \partial \Xi_j / \partial x_i$.

$(\xi_1, \xi_2, \xi_3) \equiv (\Xi_1, \Xi_2, \Xi_3) - (x_1, x_2, x_3)$ in AM78 corresponds to $(0, 0, z''')$ in the present study. Noting that $\xi_j = \Xi_j - x_j$ and using the notation of AM78, their (B.1)–(B.4) may be rewritten

$$\begin{aligned} \xi_{j,i} \overline{D}^L(u_j^\xi) &= \overline{D}^L(\xi_{j,i} u_j^\xi) - u_j^\xi [(\overline{D}^L \xi_j)_i - \overline{u}_{k,i}^L \xi_{j,k}^\xi] \\ &= \overline{D}^L(\xi_{j,i} u_j^\xi) - (\overline{u}_j^L + u_j^l) u_{j,i}^l + \overline{u}_{k,i}^L \xi_{j,k}^\xi, \end{aligned} \quad (\text{A1a})$$

$$u_i^{\text{gp}} \equiv -\overline{\xi_{j,i} u_j^l} = -\overline{\xi_{j,i} u_j^\xi}, \quad \text{and} \quad (\text{A1b})$$

$$-\overline{\xi_{j,i} \overline{D}^L(u_j^\xi)} = \overline{D}^L(u_i^{\text{gp}}) + \overline{u_j^l (u_{j,i}^l)} + \overline{u_{k,i}^L (u_k^{\text{gp}})}, \quad (\text{A1c})$$

where $\overline{D}^L \xi_j = u_j^l = u_j^\xi - \overline{u}_j^L$ is understood. The quantity u_i^{gp} is the generalized pseudomomentum for waves in a nonrotating frame (AM78). The first term on the rhs of (A1c) is analogous to $-\hat{D}_i(\overline{z_X'''} w''')$ in (20). The second term of (A1c) is analogous to $+w_X'' w''$ in (20). The third term of (A1c) is analogous to $-\hat{V}_X \cdot (\overline{\mathbf{V} z'''}) w''$ in (20).

APPENDIX B

The Derivation of the Vortex Force

In section 3c, the TWM momentum equations are rewritten for the development of the equivalent of the quasi-EM velocity, namely the total transport velocity minus the VL pseudomomentum vector. The last term in each of (36a)–(36c) is the legacy of the lateral Reynolds stress term, which can be expanded using the recipe of CL76, as shown below.

a. The prototype vortex force equations

To manipulate (36a)–(36c), we use several identities derived from the condition of horizontally homogeneous waves (29b). The first identity is

$$\nabla \hat{A}_2 = \nabla \overline{A}_2 = \nabla \overline{A}_2^e, \quad (\text{B1})$$

where A_2 is an arbitrary quantity at $O(\alpha^2)$, and (31a)–(31b) have been used. The second identity is that the vertical component of the quasi-Stokes velocity is zero, as shown by (32b), which leads to

$$\begin{aligned} \hat{D}_T &= \overline{D}_T^e + \mathbf{V}_2^{\text{qs}} \cdot \nabla + w_2^{\text{qs}} \partial_z \\ &= \overline{D}_T^e + \mathbf{V}_2^{\text{qs}} \cdot \nabla, \end{aligned} \quad (\text{B2})$$

where $\overline{D}_T^e \equiv \partial_T + \overline{\mathbf{V}}^e \cdot \nabla + \overline{w}^e \partial_z$ is the material derivative operator based on the EM velocity. The third identity is

$$-\overline{(\nabla z_1''') w_1'''} = \overline{z_1''' \nabla_1'''}, \quad (\text{B3})$$

which has been derived using (29b) and (34) and is the relationship found in the second term on the rhs of (38a). Using (B1)–(B3), we rewrite (36a)–(36c) as

$$\begin{aligned} \overline{D}_T^e \overline{\mathbf{V}}_2^e &= -\nabla(\overline{p}_4 + \overline{\eta}_4 + \overline{\pi}_4) - (\nabla \times \overline{\mathbf{V}}_2^e) \times \overline{z_1''' \nabla_1'''} \\ &\quad - \overline{(\nabla''' \cdot \nabla \mathbf{V}''')} \quad \text{and} \end{aligned} \quad (\text{B4a})$$

$$\begin{aligned} \overline{D}_T^e \overline{w}_2^e &= -\partial_z(\overline{p}_4 + \overline{\pi}_4) - \mathbf{V}_2^{\text{qs}} \cdot \nabla \overline{w}_2^e + \hat{V}_{2z} \cdot \overline{z_1''' \nabla_1'''} \\ &\quad + [\overline{\nabla''' \cdot (w_z''' \nabla z'''} - z_z''' \nabla w''')}]_4 - \overline{(\nabla''' \cdot \nabla w''')} \quad \text{and} \end{aligned} \quad (\text{B4b})$$

These equations contain a prototype of the vortex force.^{B1} The analysis up to this point was sufficient for Leibovich (1980) to derive the vortex force. This is because he used the three-dimensional Lagrangian framework of AM78. By contrast, we still have the last term of each of (B4a)–(B4b) to work on further because the VL framework is Eulerian in the horizontal direction. To deal with these terms, we use the fact that the lateral advection of velocity in the VL coordinates can be written as

$$\overline{\nabla''' \cdot \nabla \mathbf{V}'''} = \frac{1}{2} \overline{\nabla |\mathbf{V}'''|^2} + \overline{(\nabla \times \mathbf{V}''') \times \mathbf{V}'''} \quad \text{and} \quad (\text{B5a})$$

$$\overline{\nabla''' \cdot \nabla w'''} = \frac{1}{2} \partial_z \overline{|\mathbf{V}'''|^2} + \overline{(\nabla w''' - \mathbf{V}_z''') \cdot \mathbf{V}'''}, \quad (\text{B5b})$$

where here $\nabla \times \mathbf{V}''' = (v_x''' - u_y''') \mathbf{z}$, and manipulate the last term of (B5a)–(B5b) using the approach of CL76, as shown below.

b. Vorticity equations: Application of Craik and Leibovich (1976)

First we rewrite the momentum equations (11a)–(11b) in the VL coordinates as

^{B1} Note that in (B4a), the cross-product operator is the vector invariant cross product. This is because for low-pass-filtered quantities, the VL coordinates correspond to the standard Eulerian–Cartesian coordinates as pointed out by Jacobson and Aiki (2006).

$$\mathbf{V}_t + (\mathbf{V} \times \mathbf{V}) \times \mathbf{V} + \varpi \mathbf{V}_z = -\nabla(p + \eta + |\mathbf{V}|^2/2) - (\mathcal{D}_t w) \nabla z''' \quad \text{and} \quad (\text{B6a})$$

$$w_t + (\nabla w - \mathbf{V}_z) \cdot \mathbf{V} + \varpi w_z = -\partial_z(p + |\mathbf{V}|^2/2) - (\mathcal{D}_t w) z'''_z, \quad (\text{B6b})$$

where another version of (B5a)–(B5b) has been used to rewrite the advection terms. Because both $O(\alpha)$ and $O(\alpha^2)$ wave motions satisfy the apparent irrotational condition (34) and ϖ'' is sufficiently small, momentum equations for $O(\alpha^3)$ waves may be written

$$\mathbf{V}_{3\tau}''' + (\mathbf{V} \times \bar{\mathbf{V}}_2) \times \mathbf{V}_1''' + \hat{\omega}_2 \mathbf{V}_{1z}''' = -\nabla(p + \eta + |\mathbf{V}|^2/2)_3''' - [(\mathcal{D}_t w) \nabla z''']_3''' \quad \text{and} \quad (\text{B7a})$$

$$w_{3\tau}''' + (\nabla \bar{w}_2 - \bar{\mathbf{V}}_{2z}) \cdot \mathbf{V}_1''' + \hat{\omega}_2 w_{1z}''' = -\partial_z(p + |\mathbf{V}|^2/2)_3''' - [(\mathcal{D}_t w) z''']_3''' . \quad (\text{B7b})$$

Cross derivative of the above equations yields

$$(\mathbf{V} \times \mathbf{V}_3''')_\tau + \underbrace{\nabla \phi_{1\tau}'''}_{\cos\theta} \cdot \nabla (\mathbf{V} \times \bar{\mathbf{V}}_2) - (\mathbf{V} \times \bar{\mathbf{V}}_2) \underbrace{\phi_{1zz\tau}'''}_{\sin\theta} + \nabla \hat{\omega}_2 \times \underbrace{\nabla \phi_{1z\tau}'''}_{\cos\theta} = -\underbrace{[\nabla (\mathcal{D}_t w) \times \nabla z''']_3'''}_0, \quad (\text{B8a})$$

$$\begin{aligned} & (w_{3x}''' - u_{3z}''')_\tau + [(\mathbf{V} \times \bar{\mathbf{V}}_2) \underbrace{\phi_{1y\tau}'''}_{\cos\theta}]_z + (\nabla \bar{w}_2 - \bar{\mathbf{V}}_{2z})_x \cdot \underbrace{\nabla \phi_{1\tau}'''}_{\cos\theta} + (\nabla \bar{w}_2 - \bar{\mathbf{V}}_{2z}) \cdot \underbrace{\nabla \phi_{1x\tau}'''}_{\sin\theta} + \hat{\omega}_{2x} \underbrace{\phi_{1zz\tau}'''}_{\sin\theta} - \hat{\omega}_{2z} \underbrace{\phi_{1xz\tau}'''}_{\cos\theta} \\ & = \underbrace{[(\mathcal{D}_t w)_z z''']_x - (\mathcal{D}_t w)_x z''']_z}_3''' , \quad \text{and} \quad (\text{B8b}) \\ & \qquad \qquad \qquad \sin\theta \end{aligned}$$

$$\begin{aligned} & (w_{3y}''' - v_{3z}''')_\tau - [(\mathbf{V} \times \bar{\mathbf{V}}_2) \underbrace{\phi_{1x\tau}'''}_{\cos\theta}]_z + (\nabla \bar{w}_2 - \bar{\mathbf{V}}_{2z})_y \cdot \underbrace{\nabla \phi_{1\tau}'''}_{\cos\theta} + (\nabla \bar{w}_2 - \bar{\mathbf{V}}_{2z}) \cdot \underbrace{\nabla \phi_{1y\tau}'''}_{\sin\theta} + \hat{\omega}_{2y} \underbrace{\phi_{1zz\tau}'''}_{\sin\theta} - \hat{\omega}_{2z} \underbrace{\phi_{1yz\tau}'''}_{\cos\theta} \\ & = \underbrace{[(\mathcal{D}_t w)_z z''']_y - (\mathcal{D}_t w)_y z''']_z}_3''' , \quad (\text{B8c}) \\ & \qquad \qquad \qquad \sin\theta \end{aligned}$$

where $(\mathbf{V}_1''', w_1''') = (\nabla \phi_{1\tau}''', \phi_{1z\tau}''')$ and $\nabla^2 \phi_1''' + \phi_{1zz}''' = 0$ have been used. The rhs of (B8a) vanishes because both $O(\alpha)$ and $O(\alpha)$ waves are proportional to $\nabla\theta = (k, l)$ (a similar discussion appears in footnote 8).

We now use again the fact that $O(\alpha)$ and $O(\alpha^2)$ waves satisfy the apparent irrotational condition (34) to write the last term of (B5a)–(B5b) at $O(\alpha^4)$ as

$$\begin{aligned} \overline{[(\mathbf{V} \times \mathbf{V}''') \times \mathbf{V}''']}_4 &= \overline{(\mathbf{V} \times \mathbf{V}_3''') \times \mathbf{V}_1'''} \\ &= -\overline{(\mathbf{V} \times \mathbf{V}_3''')_\tau \times \nabla \phi_1'''} \\ &= -(\mathbf{V} \times \bar{\mathbf{V}}_2) \times \overline{\phi_{1zz\tau}'''} \nabla \phi_1'' \\ &= (\mathbf{V} \times \bar{\mathbf{V}}_2^e) \times \overline{z_{1z}'''} \nabla_1''', \quad (\text{B9a}) \end{aligned}$$

where the third line has been derived by first substituting for $(\mathbf{V} \times \mathbf{V}_3''')_\tau$ using (B8a) and then retaining terms labeled $\sin\theta$ in (B8a) because $\nabla \phi_1'''$ is proportional to $\sin\theta$, and

$$\begin{aligned} & \overline{[(\nabla w''' - \mathbf{V}_z''') \cdot \mathbf{V}''']}_4 \\ &= \overline{(\nabla w_3''' - \mathbf{V}_{3z}''') \cdot \mathbf{V}_1'''} \\ &= -\overline{(\nabla w_3''' - \mathbf{V}_{3z}''')_\tau \cdot \nabla \phi_1'''} \\ &= -(\nabla \bar{w}_2 - \bar{\mathbf{V}}_{2z}) \cdot \overline{(\nabla \phi_{1\tau}''')(\phi_{1xx}'' + \phi_{1yy}'')} \\ &\quad + \nabla \hat{\omega}_2 \cdot \overline{(\nabla \phi_1''') \phi_{1zz\tau}'''} \\ &\quad - \overline{[(\mathcal{D}_t w)_z \nabla z'' - z_z''' \nabla (\mathcal{D}_t w)]}_3 \cdot \nabla \phi_1'' \\ &= -\bar{\mathbf{V}}_{2z} \cdot \overline{z_{1z}'''} \nabla_1''' - \overline{[(\mathcal{D}_t w)_z \nabla z'' - z_z''' \nabla (\mathcal{D}_t w)]}_3 \cdot \nabla \phi_1''', \quad (\text{B9b}) \end{aligned}$$

where the fourth line has been derived by first substituting for $(\nabla w_3''' - \mathbf{V}_{3z}''')_\tau$ using (B8b)–(B8c) and then retaining terms labeled $\sin\theta$ in (B8b)–(B8c) because $\nabla \phi_1'''$ is proportional to $\sin\theta$. The above procedure is

based on CL76. The last line of (B9b) has been derived using $\nabla \bar{w}_2 = \nabla \hat{w}_2$ which follows from (17) and (B1).

Substitution of (B9a)–(B9b) to (B4a)–(B4b), using (B5a)–(B5b), then yields

$$\overline{\mathcal{D}_T^e \bar{\mathbf{V}}_2^e} = -\nabla(\bar{p}_4 + \bar{\eta}_4 + \bar{\Pi}_4) - (\mathbf{V} \times \bar{\mathbf{V}}_2^e) \times \mathbf{V}_2^{\text{qs}} \quad \text{and} \quad (\text{B10a})$$

$$\begin{aligned} \overline{\mathcal{D}_T^e \bar{w}_2^e} = & -\partial_z(\bar{p}_4 + \bar{\Pi}_4) - (\nabla \bar{w}_2^e - \bar{\mathbf{V}}_{2z}^e) \cdot \mathbf{V}_2^{\text{qs}} + \frac{1}{2} \partial_z (|\mathbf{V}_2^{\text{qs}}|^2 - |z_{1z}''' \mathbf{V}_1'''|^2) + [\overline{\mathbf{V}'' \cdot (w_z''' \nabla z''' - z_z''' \nabla w''')}]_4 \\ & + \overline{\nabla \phi_1''' \cdot [(D_t w)''' \nabla z''' - z_z''' \nabla (D_t w)''']}]_3, \end{aligned} \quad (\text{B10b})$$

where the third last term of (B10b) has been derived using (31a)–(32a), and

$$\bar{\Pi}_4 \equiv \frac{1}{2} (\overline{|\mathbf{V}'''|^2 - w'''^2})_4 + [\overline{(\mathbf{V}'' \cdot \nabla z''') w'''}]_4, \quad (\text{B10c})$$

is the revised Bernoulli head.^{B2} The last three terms of (B10b) consist of waves up to $O(\alpha^2)$. Because $O(\alpha)$ and $O(\alpha^2)$ waves are horizontally homogeneous, these three terms can be absorbed to the Bernoulli head in (B10c) without affecting the horizontal component of wave-averaged momentum equation (B10a). Looking at (B10a)–(B10c), $O(\alpha^3)$ waves appear only in the first term of the Bernoulli head (B10c).

APPENDIX C

Discussion on the Different Scaling for the Variation of Circulations

The utility of the velocity-based expression of the form stress term (as well as the pseudomomentum in the VL framework) is not limited to the scaling of LCs in CL76. There are various choices for the scaling of the temporal and horizontal variations of circulation, as argued in Lane et al. (2007) and AG13, and these are briefly explained in this section using a generalized expression for the scaling. Let us consider circulations whose time development is m orders (in terms of α , where $m = 0, 1, 2, \dots$) slower than the phase cycle of waves, and the horizontal scale of circulations is n orders (in terms of α , where $n = 0, 1, 2, \dots$) larger than a wavelength. The time derivative operator may be decomposed as $\partial_t = \partial_\tau + \alpha^m \partial_T$ where ∂_τ operates on wave quantities and ∂_T operates on the low-pass-filtered quantities (i.e., circulations as well as the slow time evolution of the wave quantities). Likewise the lateral gradient operator may be decomposed as $\nabla = \hat{\nabla} + \alpha^n \bar{\nabla}$, where $\hat{\nabla}$ operates on wave quantities and $\bar{\nabla}$ operates on the

low-pass-filtered quantities (i.e., circulations as well as on the large spatial-scale variation of the wave quantities).^{C1} To summarize, $\partial_\tau \bar{A} = 0$ and $\partial_T \bar{A} \neq 0$ (this is as in section 3), likewise $\hat{\nabla} A = 0$ and $\bar{\nabla} A \neq 0$ for an arbitrary quantity A , leading to

$$\partial_\tau (\overline{A_1''' B_1'''}) = 0, \quad \partial_T (\overline{A_1''' B_1'''}) \neq 0, \quad \text{and} \quad (\text{C1a})$$

$$\hat{\nabla} (\overline{A_1''' B_1'''}) = 0, \quad \bar{\nabla} (\overline{A_1''' B_1'''}) \neq 0, \quad (\text{C1b})$$

where A_1''' and B_1''' are arbitrary quantities at $O(\alpha)$.

The set of wave-averaged momentum equations that contain the vortex force may be derived if $m = n + 2$. Substitution of $n = 0$ recovers the scaling for the low-pass-filtered flow used in section 3 and is the same scaling as used in CL76. Substitution of $n = 2$ recovers the scaling of circulation in an inner-coastal shelf region in McWilliams et al. (2004). The TWM system (6a)–(6c) is written as

$$\bar{\nabla} \cdot \hat{\mathbf{V}}_2 + \partial_z \hat{w}_{n+2} = 0, \quad (\text{C2a})$$

$$\begin{aligned} (\partial_T + \hat{\mathbf{V}}_2 \cdot \bar{\nabla} + \hat{w}_{n+2} \partial_z) \hat{\mathbf{V}}_2 + \mathcal{R} \mathcal{S}_{n+4}^{\mathbf{V}} \\ = -\bar{\nabla}(\bar{p}_4 + \bar{\eta}_4) + \mathcal{F} \mathcal{S}_{n+4}^{\mathbf{V}}, \quad \text{and} \end{aligned} \quad (\text{C2b})$$

$$\underbrace{(\partial_T + \hat{\mathbf{V}}_2 \cdot \bar{\nabla} + \hat{w}_2 \partial_z) \hat{w}_2 + \mathcal{R} \mathcal{S}_4^w}_{\text{present only when } n=0} = -\bar{p}_{4z}, \quad (\text{C2c})$$

where the material derivative term in (C2c) is present only when $n = 0$. The horizontal and vertical momentum equations have been written at $O(\alpha^{n+4})$ and $O(\alpha^4)$, respectively. Using the recipe of the present study, the Reynolds stress term and the form stress term in the

^{B2} $(\hat{\mathbf{V}}_2 - \bar{\mathbf{V}}_2^e)_z \cdot (z_{1z}''' \mathbf{V}_1''') + (\bar{\mathbf{V}}_2 - \bar{\mathbf{V}}_2^e)_z \cdot (z_{1z}''' \mathbf{V}_1''') = \mathbf{V}_{2z}^{\text{qs}} \cdot \mathbf{V}_2^{\text{qs}} - (z_{1z}''' \mathbf{V}_1''')_z \cdot (z_{1z}''' \mathbf{V}_1''')$.

^{C1} Note that $\partial_t A''' = \partial_\tau A''' + \alpha^m \partial_T A'''$ and $\partial_t \bar{A} = \alpha^m \partial_T \bar{A}$ for arbitrary wave and mean quantities A''' and \bar{A} . Likewise $\nabla A''' = \hat{\nabla} A''' + \alpha^n \bar{\nabla} A'''$ and $\nabla \bar{A} = \alpha^n \bar{\nabla} \bar{A}$. It should be also noted that the amplitude, wavenumber, and frequency of $O(\alpha)$ waves are constant on the time and horizontal scales of waves (i.e., $\partial_\tau A = 0$ and $\hat{\nabla} A = 0$ for $A = \mathcal{A}, k, l, \sigma$) but may vary on the time and horizontal scales of low-pass-filtered quantities (i.e., $\partial_T A \neq 0$ and $\bar{\nabla} A \neq 0$ for $A = \mathcal{A}, k, l, \sigma$). These rules and notations are the same as that in AG13.

horizontal momentum equation (C2b) can be transformed to the horizontal component of the vortex force, $\mathbf{V}_2^{\text{qs}} \times (\bar{\mathbf{V}} \times \bar{\mathbf{V}}_2^e)$, or its variant which is $O(\alpha^{n+4})$. Likewise the Reynolds stress term in the vertical momentum equation (C2c) can be transformed into the vertical component of the vortex force $\mathbf{V}_2^{\text{qs}} \cdot (\partial_z \bar{\mathbf{V}}_2^e - \bar{\mathbf{V}} \bar{w}_2^e)$ or its variant which is $O(\alpha^4)$. Another consequence is that the TWM momentum equations (C2b)–(C2c) are to be rewritten for the development of the EM velocity.

An alternative and classical form of the wave-averaged momentum equations includes the so-called radiation stress (e.g., Longuet-Higgins and Stewart 1964, hereafter LHS64; Bühler and Jacobson 2001; Mellor 2003). The depth-integrated radiation stress of LHS64 has been written by these authors as the product of $O(\alpha)$ wave quantities. The sum of the Reynolds stress term (7) and (the negative of) the form stress term (8) then appears at $O(\alpha^{n+2})$ and when integrated over the water depth reads

$$\begin{aligned} \int_{-\infty}^{\bar{\eta}} (\mathcal{R}\mathcal{S}_{n+2}^{\mathbf{V}} - \mathcal{F}\mathcal{S}_{n+2}^{\mathbf{V}}) dz &= \bar{\mathbf{V}} \cdot \int_{-\infty}^{\bar{\eta}} \overline{\mathbf{V}_1^{\text{'''}}} \overline{\mathbf{V}_1^{\text{'''}}} dz + \frac{1}{2} \bar{\mathbf{V}} \overline{\eta_1^{\text{'''2}}} - \bar{\mathbf{V}} \int_{-\infty}^{\bar{\eta}} \overline{(z_1^{\text{'''}}} p_{1z}^{\text{'''}}) dz \\ &= \bar{\mathbf{V}} \cdot \underbrace{\int_{-\infty}^{\bar{\eta}} \overline{\mathbf{V}_1^{\text{'''}}} \overline{\mathbf{V}_1^{\text{'''}}} dz}_{S_{xx}^{(1)}} + \bar{\mathbf{V}} \underbrace{\int_{-\infty}^{\bar{\eta}} \overline{(-w_1^{\text{'''2}})} dz}_{S_{xx}^{(2)}} + \bar{\mathbf{V}} \underbrace{\frac{1}{2} \overline{\eta_1^{\text{'''2}}}}_{S_{xx}^{(3)}}, \end{aligned} \quad (\text{C3})$$

where $\overline{z_1^{\text{'''}}} p_{1z}^{\text{'''}} = -\overline{z_1^{\text{'''}}} w_{1r}^{\text{'''}} = \overline{z_{1r}^{\text{'''}}} w_1^{\text{'''}}$ has been used following (C1a), and $S_{xx}^{(1)}$, $S_{xx}^{(2)}$, and $S_{xx}^{(3)}$ are the notation in LHS64. The stress terms, $\mathcal{R}\mathcal{S}_{n+2}^{\mathbf{V}}$ and $\mathcal{F}\mathcal{S}_{n+2}^{\mathbf{V}}$, are part of the horizontal component of the TWM momentum equations written at $O(\alpha^{n+2})$. In order for the tendency term of the wave-averaged momentum equations to be written at $O(\alpha^{n+2})$, the time derivative operator needs to be decomposed as $\partial_t = \partial_\tau + \alpha^n \partial_T$, which means that $m = n$. Namely, LHS64 consider circulations whose time development is n orders slower (in term of α) than the phase cycle of the waves, which is two orders faster than that in the previous paragraph (i.e., the vortex force regime). The TWM system (6a)–(6c) becomes

$$\bar{\mathbf{V}} \cdot \hat{\mathbf{V}}_2 + \partial_z \hat{w}_{n+2} = 0, \quad (\text{C4a})$$

$$\partial_T \hat{\mathbf{V}}_2 + \mathcal{R}\mathcal{S}_{n+2}^{\mathbf{V}} = -\bar{\mathbf{V}}(\bar{p}_2 + \bar{\eta}_2) + \mathcal{F}\mathcal{S}_{n+2}^{\mathbf{V}}, \quad \text{and} \quad (\text{C4b})$$

$$0 = -\bar{p}_{2z}. \quad (\text{C4c})$$

AG13 have specialized to the case of $n = 1$ (but the result holds for $n = 2, 3, \dots$) and show that the depth-dependent radiation stress term is rewritten as

$$\mathcal{R}\mathcal{S}_{n+2}^{\mathbf{V}} - \mathcal{F}\mathcal{S}_{n+2}^{\mathbf{V}} = -\partial_T \mathbf{V}_2^{\text{qs}} + \bar{\mathbf{V}} \frac{1}{2} (\overline{|\mathbf{V}_1^{\text{'''}}|^2} - \overline{w_1^{\text{'''2}}}), \quad (\text{C5})$$

which contains no singular treatment at the sea surface, in contrast to Mellor (2008a). Substitution of (C5) to (C4b) yields a wave-averaged momentum equation written for the development of the EM velocity. The last term of (C5) vanishes in the present study because of the use of deep water waves; however, the term has been kept in order for readers to see correspondence to $\hat{\zeta}$ in McWilliams et al. (2004), J in Smith (2006), and S^J in Arduin et al. (2008b).

The difference of the incompressibility conditions of the total transport velocity, (C2a) or (C4a), and the EM velocity, $\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}_2^e + \partial_z \bar{w}_{n+2}^e = 0$, yields

$$\bar{\mathbf{V}} \cdot \mathbf{V}_2^{\text{qs}} + \partial_z w_{n+2}^{\text{qs}} = 0, \quad (\text{C6})$$

which indicates that, in the presence of the slow horizontal variations of waves, the vertical component of the quasi-Stokes velocity is nonzero and scaled at $O(\alpha^{n+2})$ (cf. Tamura et al. 2012).

APPENDIX D

Viscosity Term in the Energy Equation

The last line of (47) is the product between $-(\hat{\mathbf{V}}\theta)/\sigma$ and the viscosity term of the depth-dependent energy equation as we now show. As noted by Phillips (1977), Weber et al. (2006), and AG12, the viscosity term of the depth-dependent energy equation may be separated into two terms: one is the vertical divergence of a viscosity-induced flux (noted as FluxDiv) and one is dissipation at depths excluding the thin viscous boundary layers at the sea surface (noted as Dissipation). These two terms can be obtained by substituting (43a)–(43c) into the FluxDiv–Dissipation part of (47) to give

$$\begin{aligned} \text{FluxDiv} &= [\nu_4 \overline{\mathbf{V}_1^{\text{'''}}} \cdot (\hat{\mathbf{V}} w_{1z}^{\text{'''}} + \mathbf{V}_1^{\text{'''}}) + 2\nu_4 \overline{w_1^{\text{'''}}} w_{1z}^{\text{'''}}]_z \\ &= [\nu_4 (\overline{|\mathbf{V}_1^{\text{'''}}|^2} + \overline{w_1^{\text{'''2}}})_z]_z \\ &= \sigma^2 [\nu_4 (\kappa^2 \overline{\phi_1^{\text{'''2}}} + \overline{\phi_{1z\theta}^{\text{'''2}}})_z]_z \\ &= \sigma^2 [\nu_4 (\kappa^2 \overline{\phi_1^{\text{'''2}}} + \overline{\phi_{1z}^{\text{'''2}}})_z]_z \\ &= \sigma^2 [\nu_4 (\overline{\phi_1^{\text{'''}}} \overline{\phi_{1z}^{\text{'''}}})_{zz}]_z \quad \text{and} \end{aligned} \quad (\text{D1a})$$

$$\begin{aligned}
\text{Dissipation} &= 2\nu_4 \overline{[u_{1x}''^2 + v_{1y}''^2 + w_{1z}''^2]} + \nu_4 \overline{[v_{1x}''' + u_{1y}''']^2} \\
&\quad + \overline{[w_{1x}''' + u_{1z}''']^2} + \overline{[w_{1y}''' + v_{1z}''']^2}] \\
&= 2\nu_4 \overline{[u_{1x}''^2 + v_{1y}''^2 + w_{1z}''^2]} \\
&\quad + 4\nu_4 \overline{[u_{1y}''^2 + v_{1z}''^2 + w_{1x}''^2]} \\
&= 2\nu_4 \sigma^2 \overline{[(k^4 + l^4)\phi_{1\theta}''^2 + \kappa^4 \phi_{1\theta}''^2]} \\
&\quad + 4\nu_4 \sigma^2 \overline{[(kl)^2 \phi_{1\theta}''^2 + \kappa^2 \phi_{1z}''^2]} \\
&= 4\kappa^2 \nu_4 \sigma^2 \overline{[\kappa^2 \phi_{1\theta}''^2 + \phi_{1z}''^2]} \\
&= \nu_4 \sigma^2 \overline{(\phi_1'' \phi_{1z}''')_{zzz}}, \tag{D1b}
\end{aligned}$$

where the second line of each equation has been derived using (34), and the third line of each equation has been derived using (30c)–(30d).^{D1} It should be noted that to connect to (49), use is made of (32a).

The above two terms are part of the depth-dependent wave energy equation, which may be derived taking the sum of (A6a) and (A6b) of AG12 and then picking-up $O(\alpha^6)$ terms to yield

$$\begin{aligned}
&\frac{1}{2} \partial_T (\overline{|\mathbf{V}_1'''|^2} + w_1''^2) + \partial_z [z_{1T}''' (p_1''' + \eta_1''')] \\
&\quad + \overline{\mathbf{V}} \cdot [\overline{\mathbf{V}_1''' (p_1''' + \eta_1''')}] = -\partial_z [z_\tau''' (p''' + \eta''')]_6 \\
&\quad + \text{FluxDiv} - \text{Dissipation}, \tag{D2}
\end{aligned}$$

which indicates that the vertical flux of energy is given by the first two terms on the rhs, one is induced by pressure and one is induced by turbulent viscosity. Thus, if we consider the vertical integral of (D2) from $z = -\infty$ to $z = \bar{\eta} - \delta$ (where δ is the thickness of the TVSBL), the source of wave energy is given by the combined vertical flux $-\overline{[z_\tau''' (p''' + \eta''')]_6} + \sigma^2 \nu_4 \overline{(\phi_1'' \phi_{1z}''')_{zz}}$ evaluated at $z = \bar{\eta} - \delta$. When there is no wind forcing, we obtain $\overline{[z_\tau''' (p''' + \eta''')]_6} = \sigma^2 \nu_4 \overline{(\phi_1'' \phi_{1z}''')_{zz}}$ at $z = \bar{\eta} - \delta$, with a

^{D1} The expression of the dissipation rate as given by the last line of (D1b) is associated with only irrotational wave motions in the vertical plane, in particular at depths below the base of the TVSBL. In the present study (section 4), the coefficient of turbulent viscosity (or $\nu\kappa^2\sigma$) has been scaled at $O(\alpha^4)$ and thus the dissipation rate inside the TVSBL can be neglected, which follows Phillips (1977, p. 52) and appendix B of AG12. However, if we consider breaking waves under high wind conditions, the coefficient of turbulent viscosity (or $\nu\kappa^2\sigma$) might be scaled at $O(\alpha^2)$ or $O(\alpha^3)$ in the vicinity of the sea surface (Drazen et al. 2008; Tian et al. 2010). Then the dissipation rate associated with rotational wave motions inside the TVSBL (which is no longer thin) should be retained by revisiting (43a)–(43c) and (D1a)–(D1b), a topic we shall discuss in a later paper.

consequence that the vertical integral of the first two terms on the rhs of (D2) cancel each other. Wave energy is then gradually decreased by the last term of (D2). Nevertheless the viscosity-induced momentum flux $\nu_4 \mathbf{V}_{2z}^{\text{qs}}$ at $z = \bar{\eta} - \delta$ in (49) is nonzero, so that it remains to control the boundary condition of the EM velocity in (49).

REFERENCES

- Aiki, H., and K. J. Richards, 2008: Energetics of the global ocean: The role of layer-thickness form drag. *J. Phys. Oceanogr.*, **38**, 1845–1869.
- , and R. J. Greatbatch, 2012: Thickness-weighted mean theory for the effect of surface gravity waves on mean flows in the upper ocean. *J. Phys. Oceanogr.*, **42**, 725–747.
- , and —, 2013: The vertical structure of the surface wave radiation stress on circulation over a sloping bottom as given by thickness-weighted-mean theory. *J. Phys. Oceanogr.*, **43**, 149–164.
- Andrews, D. G., 1983: A finite-amplitude Eliassen–Palm theorem in isentropic coordinates. *J. Atmos. Sci.*, **40**, 1877–1883.
- , and M. E. McIntyre, 1978: An exact theory of nonlinear waves on a Lagrangian-mean flow. *J. Fluid Mech.*, **89**, 609–646.
- Ardhuin, F., A. D. Jenkins, and K. A. Belibassakis, 2008a: Comments on “The three-dimensional current and surface wave equations” by George Mellor. *J. Phys. Oceanogr.*, **38**, 1340–1350.
- , N. Raschle, and K. Belibassakis, 2008b: Explicit wave-averaged primitive equations using a generalized Lagrangian mean. *Ocean Modell.*, **20**, 35–60.
- Bleck, R., 1985: On the conversion between mean and eddy components of potential and kinetic energy in isentropic and isopycnic coordinates. *Dyn. Atmos. Oceans*, **9**, 17–37.
- Broström, G., K. H. Christensen, and J. E. H. Weber, 2008: A quasi-Eulerian, quasi-Lagrangian view of surface-wave-induced flow in the ocean. *J. Phys. Oceanogr.*, **38**, 1122–1130.
- Bühler, O., and T. Jacobson, 2001: Wave-driven currents and vortex dynamics on barred beaches. *J. Fluid Mech.*, **449**, 313–339.
- Chang, M.-S., 1969: Mass transport in deep-water long-crested random-gravity waves. *J. Geophys. Res.*, **6**, 1515–1536.
- Craik, A. D. D., 1985: *Wave Interactions and Fluid Flows*. Cambridge University Press, 322 pp.
- , and S. Leibovich, 1976: A rational model for Langmuir circulations. *J. Fluid Mech.*, **73**, 401–426.
- de Szoeke, R. A., and A. F. Bennett, 1993: Microstructure fluxes across density surfaces. *J. Phys. Oceanogr.*, **23**, 2254–2264.
- Dingemans, M. W., 2009: Some reflection on the generalized Lagrangian mean. *Nonlinear Wave Dynamics*, World Scientific Publishing, 1–30.
- Donelan, M. A., M. Curcic, S. S. Chen, and A. K. Magnusson, 2012: Modeling waves and wind stress. *J. Geophys. Res.*, **117**, C00J23, doi:10.1029/2011JC007787.
- Drazen, D. A., W. K. Melville, and L. Lenain, 2008: Inertial scaling of dissipation in unsteady breaking waves. *J. Fluid Mech.*, **611**, 307–332.
- Fan, Y., I. Ginis, and T. Hara, 2010: Momentum flux budget across the air–sea interface under uniform and tropical cyclone winds. *J. Phys. Oceanogr.*, **40**, 2221–2242.
- Gallimore, R. G., and D. R. Johnson, 1981: The forcing of the meridional circulation of the isentropic zonally averaged circumpolar vortex. *J. Atmos. Sci.*, **38**, 583–599.

- Garrett, C., 1976: Generation of Langmuir circulations by surface waves—A feedback mechanism. *J. Mar. Res.*, **34**, 117–130.
- Greatbatch, R. J., 1998: Exploring the relationship between eddy-induced transport velocity, vertical momentum transfer, and the isopycnal flux of potential vorticity. *J. Phys. Oceanogr.*, **28**, 422–432.
- , and T. J. McDougall, 2003: The non-Boussinesq temporal residual mean. *J. Phys. Oceanogr.*, **33**, 1231–1239.
- Hasselmann, K., 1970: Wave-driven inertial oscillations. *Geophys. Astrophys. Fluid Dyn.*, **1**, 463–502.
- Huang, N. E., 1979: On surface drift currents in the ocean. *J. Fluid Mech.*, **91**, 191–208.
- Iwasaki, T., 2001: Atmospheric energy cycle viewed from wave-mean-flow interaction and Lagrangian mean circulation. *J. Atmos. Sci.*, **58**, 3036–3052.
- Jacobson, T., and H. Aiki, 2006: An exact energy for TRM theory. *J. Phys. Oceanogr.*, **36**, 558–564.
- Jenkins, A. D., 1987: Wind- and wave-induced currents in a rotating sea with depth-varying eddy viscosity. *J. Phys. Oceanogr.*, **17**, 938–951.
- , 1989: The use of a wave prediction model for driving a near-surface current model. *Dtsch. Hydrogr. Z.*, **42**, 133–149.
- , and F. Ardhuin, 2004: Interaction of ocean waves and currents: How different approaches may be reconciled. *Proc. 14th Int. Offshore and Polar Engineering Conf.*, Toulon, France, International Society of Offshore and Polar Engineers, 105–111.
- Kukulka, T., A. J. Plueddemann, J. H. Trowbridge, and P. P. Sullivan, 2010: Rapid mixed layer deepening by the combination of Langmuir and shear instabilities: A case study. *J. Phys. Oceanogr.*, **40**, 2381–2400.
- Lagrange, J. L., 1788: *Mécanique Analytique*. La Veuve Desaint, 512 pp.
- Lamb, H., 1932: *Hydrodynamics*. 6th ed. Cambridge University Press, 738 pp.
- Lane, E. M., J. M. Restrepo, and J. C. McWilliams, 2007: Wave-current interaction: A comparison of radiation-stress and vortex-force representations. *J. Phys. Oceanogr.*, **37**, 1122–1141.
- Langmuir, I., 1938: Surface motion of water induced by wind. *Science*, **87**, 119–123.
- Large, W. G., and S. Pond, 1981: Open ocean momentum flux measurements in moderate to strong winds. *J. Phys. Oceanogr.*, **11**, 324–336.
- Leibovich, S., 1980: On wave-current interaction theory of Langmuir circulations. *J. Fluid Mech.*, **99**, 715–724.
- Longuet-Higgins, M. S., 1953: Mass transport in water waves. *Phil. Trans. Roy. Soc. London*, **A245**, 535–581.
- , 1960: Mass transport in the boundary layer at a free oscillating surface. *J. Fluid Mech.*, **8**, 293–306.
- , and R. W. Stewart, 1964: Radiation stress in water waves: A physical discussion with applications. *Deep-Sea Res.*, **11**, 529–562.
- McDougall, T. J., and P. C. McIntosh, 2001: The temporal-residual-mean velocity. Part II: Isopycnal interpretation and the tracer and momentum equations. *J. Phys. Oceanogr.*, **31**, 1222–1246.
- McWilliams, J. C., J. M. Restrepo, and E. M. Lane, 2004: An asymptotic theory for the interaction of waves and currents in coastal waters. *J. Fluid Mech.*, **511**, 135–178.
- Mellor, G., 2003: The three-dimensional current and surface wave equations. *J. Phys. Oceanogr.*, **33**, 1978–1989.
- , 2008a: The depth-dependent current and wave interaction equations: A revision. *J. Phys. Oceanogr.*, **38**, 2587–2596.
- , 2008b: Reply. *J. Phys. Oceanogr.*, **38**, 1351–1353.
- Ng, C.-O., 2004: Mass transport in gravity waves revisited. *J. Geophys. Res.*, **109**, C04012, doi:10.1029/2003JC002121.
- Phillips, O. M., 1977: *Dynamics of the Upper Ocean*. Cambridge University Press, 336 pp.
- Piedra-Cueva, I., 1995: Drift velocity of spatially decaying waves in a two-layer viscous system. *J. Fluid Mech.*, **299**, 217–239.
- Pierson, W. J., 1962: Perturbation analysis of the Navier–Stokes equations in Lagrangian form with selected linear solutions. *J. Geophys. Res.*, **67**, 3151–3160.
- Polton, J. A., and S. E. Belcher, 2007: Langmuir turbulence and deeply penetrating jets in an unstratified mixed layer. *J. Geophys. Res.*, **112**, C09020, doi:10.1029/2007JC004205.
- Skyllingstad, E. D., and D. W. Denbo, 1995: An ocean large-eddy simulation of Langmuir circulations and convection in the surface mixed layer. *J. Geophys. Res.*, **100** (C5), 8501–8522.
- Smith, J. A., 2006: Wave-current interactions in finite depth. *J. Phys. Oceanogr.*, **36**, 1403–1419.
- Stokes, G. G., 1847: On the theory of oscillatory waves. *Trans. Cambridge Philos. Soc.*, **8**, 441–455.
- Tamura, H., Y. Miyazawa, and L.-Y. Oey, 2012: The Stokes drift and wave induced-mass flux in the North Pacific. *J. Geophys. Res.*, **117**, C08021, doi:10.1029/2012JC008113.
- Tang, C. L., W. Perrie, A. D. Jenkins, B. M. DeTracey, Y. Hu, B. Toulany, and P. C. Smith, 2007: Observation and modeling of surface currents on the Grand Banks: A study of the wave effects on surface currents. *J. Geophys. Res.*, **112**, C10025, doi:10.1029/2006JC004028.
- Tian, Z., M. Perlin, and W. Choi, 2010: Energy dissipation in two-dimensional unsteady plunging breakers and an eddy viscosity model. *J. Fluid Mech.*, **655**, 217–257.
- Ünlüata, U., and C. C. Mei, 1970: Mass transport in water waves. *J. Geophys. Res.*, **75**, 7611–7618.
- Weber, J. E., 1983: Steady wind- and wave-induced currents in the open ocean. *J. Phys. Oceanogr.*, **13**, 524–530.
- , G. Broström, and O. Saetra, 2006: Eulerian versus Lagrangian approaches to the wave-induced transport in the upper ocean. *J. Phys. Oceanogr.*, **36**, 2106–2118.
- Xu, Z., and A. J. Bowen, 1994: Wave- and wind-driven flow in water of finite depth. *J. Phys. Oceanogr.*, **24**, 1850–1866.
- Young, W. R., 2012: An exact thickness-weighted average formulation of the Boussinesq equations. *J. Phys. Oceanogr.*, **42**, 692–707.