

Bibliography (papers published by IFREE researchers)

- Aizawa, Y., K. Ito, and Y. Tatsumi, Experimental determination of compressional wave velocities of olivine aggregate up to 1000°C at 1 GPa, *Tectonophysics*, 339, 473-478, 2001.
- Baba, T., T. Hori, S. Hirano, P. R. Cummins, J. -O. Park, M. Kameyama, and Y. Kaneda, Deformation of a seamount subducting beneath an accretionary prism: constraints from numerical simulation, *Geophys. Res. Lett.*, 28, 1827-1830, 2001.
- Baba, T., and Y. Tanioka, The shape of Philippine Sea Plate estimated from seismic surveys and the slip distribution of the 1946 Nankai earthquake (in Japanese), *Monthly Earth*, 23, 697-702, 2001
- Baba, T., Y. Tanioka, P. R. Cummins, and K. Uhira, The slip distribution of the 1946 Nankai earthquake estimated from tsunami inversion using a new plate model, *Phys. Earth Planet. Int.*, 132, 59-73, 2002.
- Borissova, I., A. Moore, J. Sayers, R. Parums, M. F. Coffin, and P. A. Symonds, Geological Framework of the Kerguelen Plateau and Adjacent Ocean Basins, *Record 2002/05, Geoscience Australia*, 120pp., 2002.
- Coffin, M. F., Kerguelen Plateau, and Broken Ridge, in J. Nybakken, ed., *Encyclopedia of Oceanography and Marine Science*, Fitzroy Dearborn, Chicago, in press.
- Coffin, M. F., M. S. Pringle, R. A. Duncan, T. P. Gladzhenko, M. Storey, R. D. Müller, and L. A. Gahagan, Kerguelen hotspot magma output since 130 Ma, *J. Petrol.*, 43, 1121-1139, 2002.
- Cummins, P. R., T. Baba, T. Hori, and Y. Kaneda, The influence of Philippine Sea Plate structure on great Nankai Trough earthquakes inferred from the rupture process of the 1946 Nankai earthquake, *J. Geography*, 110, 498-509, 2001.
- Cummins, P. R., T. Hori, and Y. Kaneda, Splay fault and megathrust earthquake slip in the Nankai Trough, *Earth Planet Space*, 53, 243-248, 2001.
- Cummins, P. R., T. Baba, S. Kodaira, and Y. Kaneda, The 1946 Nankai earthquake and segmentation of the Nankai Trough, *Phys. Earth Planet. Int.*, 132, 75-87, 2002.
- Frey, F. A., M. F. Coffin, P. J. Wallace, and P. Quilty, eds., *Proceedings of the Ocean Drilling Program, Scientific Results*, 183, College Station TX (Ocean Drilling Program), in press.
- Frey, F. A., M. F. Coffin, P. J. Wallace, and D. Weis, Leg 183 synthesis: Kerguelen Plateau-Broken Ridge - a large igneous province, in F. A. Frey, M. F. Coffin, P. J. Wallace, and P. Quilty, eds., *Proc. ODP, Sci. Res.*, 183, College Station TX (Ocean Drilling Program), in press.
- Fukao, Y., S. Widiyantoro, and M. Obayashi, Stagnant slabs in the upper and lower mantle transition zone, *Rev. Geophys.*, 39, 291-323, 2001.
- Fukuyama, E., C. Hashimoto, and M. Matsu'ura, Simulation of the transition of earthquake rupture from quasi-static growth to dynamic propagation, *Pure Appl. Geophys.*, 159, 2057-2066, 2002.
- Fukuyama, E., C. Hashimoto, and M. Matsu'ura, Simulation of earthquake rupture transition from quasi-static growth to dynamic propagation, in *2nd ACES Workshop Proc.*, M. Matsu'ura, K. Nakajima, and P. Mora, eds., APEC Cooperation for Earthquake Simulation, 375-380, 2001.
- Gooday, A. J., H. Kitazato, S. Hori, and T. Toyofuku, Monothalamous soft-shelled foraminifera at an abyssal site in the North Pacific: a preliminary report, *J. Oceanogr.*, 57, 377-384, 2001.
- Gorvatov, A., Y. Fukao, and S. Widiyantoro, Application of a three-dimensional ray-tracing technique to global P, PP and Pdiff travel time tomography, *Geophys. J. Int.*, 146, 583-593, 2001.
- Gorvatov, A., Y. Fukao, S. Widiyantoro, and E. Gordeev, Seismic evidence for a mantle plume oceanward of the Kamchatka-Aleutian trench junction, *Geophys. J. Int.*, 146, 282-288, 2001.
- Hanyu, T., Y. Tatsumi, and S. Nakai, A contribution of slab-melts to the formation of high-Mg andesite magmas; Hf isotopic evidence from SW Japan, *Geophys. Res. Lett.*, in press, 2002.
- Hasegawa, T., H. Nishi, H. Okada, T. Sakamoto, L. Beufort, F. Giraud, O. Friedrich, M. Furukawa, H. Kawahata, N. Ohkouchi, R. Takashima, A. Kuroyamagi, M. Yamamura, and N. Katsuta, High resolution analyses of late Cretaceous Ocean Anoxic Event (OAE)-1b (in Japanese), *Monthly Earth*, 24, 454-460, 2002.
- Hashimoto, C., and M. Matsu'ura, 3-D simulation of earthquake generation cycles and evolution of fault constitutive properties, *Pure Appl. Geophys.*, 159, 2175-2199, 2002.
- Hashimoto, C., and M. Matsu'ura, Physical modelling of earthquake generation cycles at transcurent plate boundaries, in *2nd ACES Workshop Proc.*, M. Matsu'ura, K. Nakajima, and P. Mora, eds., APEC Cooperation for Earthquake Simulation, 211-215, 2001.
- Hattori, Y., K. Suzuki, M. Honda, and H. Shimizu, Re-Os isotope systematics of the Taklimakan Desert sands, moraines and river sediments around the Taklimakan Desert, and of Tibetan soils, *Geochim. Cosmochim. Acta*, in press, 2002.
- Heinz, P., H. Kitazato, G. Schmiedl, and Ch. Hemleben, Response of deep-sea benthic foraminifera from the Mediterranean Sea to simulated phytoplankton pulses under laboratory conditions, *J. Foram. Res.*, 31, 210-227, 2001.
- Heinz, P., Ch. Hemleben, and H. Kitazato, Time-response of cultured deep-sea benthic foraminifera to different algal diets, *Deep Sea Res.*, I, 49, 517-537, 2002.
- Hirano, S., Y. Ogawa, and K. Kawamura, Deformation of un lithified Sediments in an early stage of the compaction process deduced from microtextures and magnetic fabrics: ODP Leg 174B, Hole 1074A, in *Proc. ODP, Sci. Res.*, K. Becker, and M. J. Malone, eds., 174B, 1-13, College Station, TX.
- Hisamitu, T., and K. Ujiie, Early deformation state of prism and underthrust sediments resulted by study of anisotropy of magnetic susceptibility (AMS) (in Japanese), *Monthly Earth*, 24, 200-207, 2001.
- Hisamitu, T., and T. Kanematsu, Changes in magnetic fabrics of sediments at the western Pacific subduction margins, *Monthly Earth*, 32, 191-199, 2001.
- Hori, T., T. Baba, P. R. Cummins, and Y. Kaneda, Effects of subducted seamounts on the source process of the 1946 Nankai Earthquake, SW Japan, *2nd ACES Workshop Proc.*, 271-274, 2001.
- Hori, T., and Y. Kaneda, A simple explanation for the occurrence of the 1911 Morgan Hill earthquake in the stress shadow of the 1906 San Francisco earthquake, *Geophys. Res. Lett.*, 28, 2261-2264, 2001.
- Hori, T., and Y. Kaneda, Stress accumulation and stress change in the critically stressed crust with a weak fault, *Proc. Int. Symp. on slip and flow processes in and below the seismogenic region*, 387-388, 2001.
- Hori, T., Transition from quiet to active period of seismicity

- around source regions of great interplate earthquakes (in Japanese with English abstract), *J. Geography*, *111*, 192-199, 2002.
- Hori, T., and K. Oike, Active and quiet periods of seismicity in Southwest Japan (in Japanese), *Monthly Earth*, *38*, 14-19, 2002.
- Hurukawa, N., N. Seto, H. Inoue, K. Nishigami, I. Marzouk, A. Megahed, E. M. Ibrahim, H. Murakami, N. Nakamura, T. Haneda, S. Sugiyama, T. Ohkura, Y. Fujii, H. M. Hussein, A. S. Megahed, H. F. Mohammed, R. A. Fattah, M. Mizoue, S. Hashimoto, M. Kobayashi, and D. Suetsugu, Seismological observations in and around the Southern part of the Gulf of Suez, Egypt, *Bull. Seismol. Soc. Am.*, *91*, 708-717, 2001.
- Ichiki, M., M. Uyeshima, H. Utada, G. Zhao, J. Tang, and M. Ma, Upper mantle conductivity structure of the back-arc region beneath northeastern China, *Geophys. Res. Lett.*, *28*, 3773-3776, 2001.
- Itou, M., T. Ono, T. Oba, and S. Noriki, Isotopic composition and morphology of living *Globorotalia scitula*: a new proxy of sub-intermediate ocean carbonate chemistry?, *Mar. Micropaleontol.*, *42*, 189-210, 2001.
- Kameyama, M., T. Hori, P. R. Cummins, S. Hirano, T. Baba, K. Uhira, and Y. Kaneda, An optimistic interpretation of friction law from thermal-mechanical coupling in shear deformation of viscoelastic material, *2nd ACES Workshop Proc.*, 91-95, 2001.
- Kameyama, M., T. Hori, P. R. Cummins, S. Hirano, T. Baba, and Y. Kaneda, Numerical experiments of shear deformation with frictional heating (in Japanese with English abstract), *Bull. Earthq. Res. Inst. Univ. Tokyo*, *75*, 127-143, 2000.
- Kameyama, M., and Y. Kaneda, Thermal-mechanical coupling in shear deformation of viscoelastic material as a model of frictional constitutive relations, *Pure Appl. Geophys.*, *159*, 2011-2028, 2002.
- Kato, A., M. Ohnaka, and H. Mochizuki, Constitutive properties for the shear failure of intact granite in seismogenic environments, *J. Geophys. Res.*, in press, 2002.
- Kido, Y., Geophysical properties of the Northern South China Sea, the continental margin -an example of utilization of Jamstec Frontier Database System, *Geoinformatics*, *11*, 223-233, 2001.
- Kido, Y., K. Suyehiro, and H. Kinoshita, Rifting and spreading of the continental margin of the South China Sea -as a case of Frontier Database study, *Mar. Geophys. Res.*, *22*, 1-15, 2001.
- Kido, Y., S. Machida, H. Sato, and K. Fujioka, Origin of magnetic dipole anomaly of Shikoku -an example of utilization of Jamstec Frontier Database System, *JAM-STECC J. Deep Sea Res.*, *18*, 171-183, 2001.
- Kido, Y., T. Tsuru, J. -O. Park, T. Higashikata, Y. Kaneda, and Y. Kono, Three dimensional overview of the Japan trench -an example of utilization of the Frontier Database System, *Computers & Geosciences*, *27*, 43-57, 2001.
- Kido, Y., Y. Kosumi, A. Nakanishi, T. Tsuru, and Y. Kaneda, Geophysical features on subducting seamounts at the junction of Japan and Kuril Trenches -Using potential anomalies and admittance functions, *Geoinformatics*, *13*, 3, 141-151, 2002.
- Kido, Y., K. Fujioka, S. Machida, and H. Sato, A serpentine diapir as a source of magnetic dipole anomalies along a tectonic line in the southwest Japan, *Geophys. Res. Lett.*, in press, 2002.
- Kitazato, H., Life cycles and life habits, in *Ecology of Extinct Organisms, Science of Extinct Organisms* (in Japanese), 3, N. Ikeya and K. Tanabe, eds., Asakura Shoten Publ., Tokyo, 2001.
- Kitazato, H., Meanings of foraminiferal test morphology: Actinopaleontological Approach (in Japanese), *Japanese Scientific Monthly*, *54*, 46-50, 2001.
- Kitazato, H., Sedimentary process of both organic and clastic materials at subduction zone: exemplified from Sagami Bay (in Japanese), *Monthly Earth, Spec.*, *32*, 74-81, 2001.
- Kitazato, H., M. Tsuchiya, and M. McGann, How does Trochammina hadai Uchio, blackish foraminifera, that dwells in East Asia, migrate beyond the Pacific Ocean? (in Japanese), *Monthly Kaiyo*, *33*, 399-405, 2001.
- Kitazato, H., New approaches to fossils as living organisms (in Japanese), in *Science Reports by Takashi Tachibana, What is it? Pursuing the fabrication case of stone implements of the Old Stone Age*, T. Tachibana, ed., Asahi Shinbunsha, Tokyo, 189-192, 2001.
- Kobayashi, K., S. Hirano, T. Arai, R. Ikeda, K. Omura, H. Sano, T. Sawaguchi, H. Tanaka, T. Tomita, N. Tomida, T. Matsuda, and A. Yamazaki, Distribution of fault rocks in the fracture zone of the Nojima Fault at a depth of 1140 m: observations from the Hirabayashi NIED drill core, *The Island Arc*, *10*, 411-421, 2001.
- Kobayashi, K., T. Arai, R. Ikeda, K. Omura, K. Shimada, H. Tanaka, T. Tomita, S. Hirano, and T. Matsuda, Textures of fault-related rocks in the 1300 m damaged zone of the Nojima Fault: examples from the NIED Hirabayashi core samples, *NIED Tech. Rep.*, *61*, 223-229, 2001.
- Kodaira, S., E. Kurashimo, J. -O. Park, N. Takahashi, A. Nakanishi, S. Miura, T. Iwasaki, N. Hirata, K. Ito, and Y. Kaneda, Rupture process of the 1946 Nankaido earthquake deduced from results of wide-angle seismic survey, *Proc., First China-Japan Workshop on Earthquake Disaster Mitigation*, 18-21, 2001.
- Kodaira, S., E. Kurashimo, J. -O. Park, N. Takahashi, A. Nakanishi, S. Miura, T. Iwasaki, N. Hirata, K. Ito, and Y. Kaneda, Structural factors controlling the rupture process of a megathrust earthquake at the Nankai trough seismogenic zone, *Geophys. J. Int.*, *149*, 815-835, 2002.
- Kodaira, S., K. Uhira, T. Tsuru, H. Sugioka, K. Suyehiro and Y. Kaneda, Seismic image and its implications for an earthquake swarm at an active volcanic region off the Miyake-jima - Kozu-shima, Japan, *Geophys. Res. Lett.*, *29*, 10, 1029/2001GL014377, 2002.
- Kodaira, S., J. -O. Park, A. Nakanishi, and Y. Kaneda, Candidates of the Nankai seismogenic zone drilling and their structures (in Japanese), *Monthly Earth, Spec. Issue*, 114-118, 2002.
- Kushiro, I., Partial melting experiments on peridotite and origin of mid-ocean ridge basalt, *Ann. Rev. Earth Planet. Sci.*, *29*, 71-107, 2001.
- Kushiro, I., and B. O. Mysen, A possible effect of melt structure on the Mg-Fe²⁺ partitioning between olivine and melt, *Geochim. Cosmochim. Acta*, *66*, 2267-2272, 2002.
- Matsuda, T., T. Arai, R. Ikeda, K. Omura, K. Shimada, H. Tanaka, T. Tomita, and S. Hirano, Material analyses of the damaged zone at 1140, 1300 and 1800 m of the Nojima Fault, based on the NIED Hirabayashi core samples, *NIED Tech. Rep.*, *61*, 183-193, 2001.
- Matsuoka, J., A. Kano, T. Oba, T. Watanabe, S. Sakai, and K. Seto, Seasonal variation of stable isotopic compositions recorded in a laminated tuff, SW-Japan, *Earth Planet. Sci. Lett.*, *192*, 31-44, 2001.
- Matsuoka, K., H. Kitazato, and R. Nomura, Changes in ecosystems and their relations to geology and paleontology: invitation to the international symposium "Application of

- Micro- and Meioorganisms to Environmental Problems, EM3" (in Japanese), *Monthly Kaiyo*, 33, 379-381, 2001.
- Miura, S., N. Takahashi, A. Nakanishi, S. Kodaira, and Y. Kaneda, Seismic experiment in the Japan Trench fore arc region (off Miyagi)-KY9905 cruise, *JAMSTEC J. Deep Sea Res.*, 18, 145-156, 2001.
- Moe, K. T., H. Tokuyama, M. Murayama, and KH 00-5 scientific party, HINDOO cruise deep-sea channel survey in the Bay of Bengal, *Bull. Geol. Soc. Japan*, 107, 2001.
- Nakajima, K., C. Hashimoto, and M. Matsu'ura, Parallel performance of the tectonic loading process model at transcurrent plate boundaries, in *2nd ACES Workshop Proc.*, M. Matsu'ura, K. Nakajima, and P. Mora, eds., APEC Cooperation for Earthquake Simulation, 295-299, 2001.
- Nakanishi, A., H. Shiobara, R. Hino, K. Mochizuki, T. Sato, J. Kasahara, N. Takahashi, K. Suyehiro, H. Tokuyama, J. Segawa, M. Shinohara, and H. Shimamura, Deep crustal structure of the eastern Nankai trough and Zenisu ridge by dense airgun -OBS seismic profiling, *Mar. Geology*, 187, 47-62, 2002.
- Nakanishi, A., S. Kodaira, J. -O. Park, and Y. Kaneda, Deformable backstop as seaward end of coseismic slip in the Nankai Trough seismogenic zone, *Earth Planet. Sci. Lett.*, 203, 255-263, 2002.
- Nakanishi, A., H. Shiobara, R. Hino, J. Kasahara, K. Suyehiro, and H. Shimamura, Crustal structure around the eastern end of coseismic rupture zone of the 1944 Tonankai earthquake, *Tectonophysics*, 354, 257-275, 2002.
- Nakanishi, A., N. Takahashi, J. -O. Park, S. Miura, S. Kodaira, Y. Kaneda, N. Hirata, T. Iwasaki, and M. Nakamura, Crustal structure across the coseismic rupture zone of the 1944 Tonankai earthquake, the central Nankai Trough seismic zone, *J. Geophys. Res.*, 107, 2-21, 2002.
- Nakanishi, A., S. Miura, K. Obana, S. Kodaira, Y. Kaneda, N. Takahashi, Wide-angle OBS survey around the Kurile Trench, the eastern Hokkaido -KY0005 cruise (in Japanese with English abstract), *JAMSTEC J. Deep Sea Res.*, 18, 157-169, 2001.
- Nakanishi, A., and Y. Kaneda, Crustal structure and seismicity of the Nankai subduction seismogenic zone -results and future plan (in Japanese), *Monthly Earth*, 23, 685-696, 2001.
- Nakanishi, I., and T. Isse, the anisotropy and the differential rotation of the inner core using seismograms observed at Syowa station, Antarctica (in Japanese), *Monthly Earth*, 35, 20-25, 2001.
- Obana, K., S. Kodaira, K. Mochizuki, and M. Shinohara, Micro-seismicity around the seaward updip limit of the 1946 Nankaido earthquake dislocation area, *Geophys. Res. Lett.*, 82, 2333-2336, 2001.
- Ohkouchi, N., T. I. Eglinton, L. D. Keigwin, and J. M. Hayes, Spatial and temporal offsets between proxy records in a sediment drift, *Science*, in press, 2002.
- Ohtaki, T., T. Saita, D. Suetsugu, K. Kanjo, and H. Takenaka, The structure of the mantle transition zone beneath subduction zones in the western Pacific (in Japanese), *Monthly Earth*, 23, 455-458, 2001.
- Omura, K., R. Ikeda, T. Arai, K. Kobayashi, K. Shimada, H. Tanaka, T. Tomita, S. Hirano, and T. Matsuda, Distribution of Fault-related rocks in the 1800 m damaged zone of the Nojima Fault, based on the NIED Hirabayashi core samples, *NIED Tech. Rep.*, 61, 173-182, 2001.
- Omura, K., R. Ikeda, T. Matsuda, T. Arai, K. Kobayashi, K. Shimada, H. Tanaka, T. Tomita, and S. Hirano, Photographs of NIED Nojima Fault drilling cores at Hirabayashi borehole, *NIED Tech. Note*, 229, 2001.
- Omura, K., R. Ikeda, Y. Iio, T. Arai, K. Kobayashi, K. Shimada, H. Tanaka, T. Tomita, S. Hirano, and T. Matsuda, Characteristics of damaged zones of the Nojima Fault, based on the NIED logging data at Hirabayashi, *NIED Tech. Rep.*, 61, 155-171, 2001.
- Ono, S., Mineralogy of Peridotite and basalt at the earth's lower-mantle pressures, *Acta Cryst.*, A58 (Supplement), C39, 2002.
- Ono, S., Phase boundary of silicon dioxide SiO₂ determined by in situ X-ray and laser heating technique, *SPRING-8 Research Frontiers 2000/2001*, 59-60, 2002.
- Ono, S., E. Ito, and T. Katsura, Mineral chemistry of subducted basaltic crust (MORB) from 25 to 37 GPa, and chemical heterogeneity of the lower mantle, *Earth Planet. Sci. Lett.*, 190, 57-63, 2001.
- Ono, S., K. Hirose, M. Isshiki, K. Mibe, and Y. Saito, Equation of state of hexagonal aluminous phase of natural composition to 63 GPa at 300 K, *Phys. Chem. Mineral.*, 29, 527-531, 2002.
- Ono, S., K. Hirose, M. Murakami, and M. Isshiki, Post-stishovite phase boundary in SiO₂ determined by in situ X-ray observations, *Earth Planet. Sci. Lett.*, 197, 187-192, 2002.
- Ono, S., K. Hirose, N. Nishiyama, and M. Ishik, Phase boundary between rutile-type and CaCl₂-type germanium dioxide determined by in situ X-ray observations, *Am. Mineral.*, 87, 99-102, 2002.
- Ono, S., K. Hirose, T. Kikegawa, and Y. Saito, The compressibility of a natural composition calcium ferrite-type aluminous phase to 70 GPa, *Phys. Earth Planet. Int.*, 131, 311-318, 2002.
- Ono, S., K. Mibe, and T. Yoshino, Aqueous fluid connectivity in pyrope aggregates: water transport into the deep mantle by a subducted oceanic crust without any hydrous minerals, *Earth Planet. Sci. Lett.*, in press, 2002.
- Ono, S., T. Suto, K. Hirose, Y. Kuwayama, T. Komabayashi, and T. Kikegawa, Equation of state of Al-bearing stishovite to 40 GPa at 300 K, *Am. Mineral.*, 87, 1486-1489, 2002.
- Park, J. -O., T. Tsuru, S. Kodaira, P. R. Cummins, and Y. Kaneda, Splay Fault branching along the Nankai subduction zone, *Science*, 297, 1157-1160, 2002.
- Park, J. -O., T. Tsuru, N. Takahashi, T. Hori, S. Kodaira, A. Nakanishi, S. Miura, and Y. Kaneda, A deep strong reflector of the Nankai accretionary wedge from multichannel seismic data: implications for underplating and interseismic shear stress release, *J. Geophys. Res.*, 107, 3-16, 2002.
- Park, J. -O., T. Tsuru, T. Hamajima, Y. Kaneda, A. Taira, S. Kuramoto, and EW9907/08 Shipboard scientific party, AVO analysis along the Nankai Trough decollement (in Japanese with English abstract), *J. Geography*, 110, 510-520, 2001.
- Park, J. -O., T. Tsuru, and Y. Kaneda, Crustal structure of the Nankai Trough accretionary wedge (in Japanese), *Monthly Earth, Spec. Issue*, 160-167, 2001.
- Sagiya, T., Sato, T., Hashimoto, C., Minami, K., and M. Matsu'ura, Viscoelastic inversion of crustal deformation data, in *2nd ACES Workshop Proc.*, M. Matsu'ura, K. Nakajima, and P. Mora, eds., APEC Cooperation for Earthquake Simulation, 439-442, 2001.
- Sakai, S., and A. Kano, Original oxygen isotopic composition of planktic foraminifers preserved in diagenetically altered Pleistocene shallow-marine carbonates, *Mar. Geol.*, 172, 197-201, 2001.
- Sakamoto, T., Sedimentary rhythms and paleoceanography (in Japanese), *Monthly Earth*, 24, 410-415, 2002.
- Suetsugu, D., A low Qscs anomaly near the South Pacific

- Superswell, *Geophys. Res. Lett.*, **28**, 391-394, 2001.
- Suetsugu, D., F. Niu, and Y. Yoshida, The structure of the mantle boundary layers beneath the South Pacific (in Japanese), *Monthly Earth*, **23**, 451-454, 2001.
- Sugioka, H., Y. Fukao, T. Okamoto, and K. Kanjo, Detection of shallowest submarine seismicity by acoustic-coupled-shear waves, *J. Geophys. Res.*, **106**, 13485-13499, 2001.
- Suzuki, K., and Y. Tatsumi, Osmium concentrations and $^{187}\text{Os}/^{188}\text{Os}$ ratios of GSJ reference samples, JB-1a, JA-2 and JP-1, *Geochem. J.*, **35**, 207-210, 2001.
- Taira, A., G. F. Moore, S. Hirano, K. Ujiie, T. Hisamitsu, and Y. Murakami, Leg 190: Nankai Trough I, *ODP News Letter (ODP Japan Office)*, **18**, 12-14, 2001.
- Tamura, Y., Noto Kongo-Kurosaki volcanic rocks in the western part of Noto Peninsula, Ishikawa Prefecture (in Japanese), in *Hokuriku no Shizen o Tazunete*, Tsukiji-shokan, 126-132, 2001.
- Tamura, Y., Y. Tatsumi, D. Zhao, Y. Kido, and H. Shukuno, Distribution of Quaternary volcanoes in the Northeast Japan arc: geologic and geophysical evidence of hot fingers in the mantle wedge, *Proc. Japan Acad.*, **77**, 135-139, 2001.
- Tamura, Y., and Y. Tatsumi, Remelting of an andesitic crust as a possible origin for rhyolitic magma in oceanic arcs: an example from the Izu-Bonin arc, *J. Petrol.*, **43**, 1029-1047, 2002.
- Tamura, Y., Y. Tatsumi, D. Zhao, Y. Kido, and H. Shukuno, Hot fingers in the mantle wedge: new insights into magma genesis in subduction zones, *Earth Planet. Sci. Lett.*, **197**, 105-116, 2002.
- Tanaka, H., T. Matsuda, K. Omura, R. Ikeda, K. Kobayashi, K. Shimada, T. Arai, T. Tomita, and S. Hirano, Complete fault rock distribution analysis along the Hirabayashi NIED Core penetrating the Nojima Fault at 1140m depth, Awaji Island, Southwest Japan, *Ret. NIED*, **61**, 195-221, 2001.
- Tatsumi, Y., Ancient activity of the South Pacific superplume (in Japanese), *Monthly Earth*, **23**, 488-492, 2001.
- Tatsumi, Y., Geochemical modeling of partial melting of subducting sediments and subsequent melt-mantle interaction: generation of high-Mg andesites in the Setouchi volcanic belt, southwest Japan, *Geology*, **29**, 323-326, 2001.
- Tatsumi, Y., N. Ishikawa, K. Anno, K. Ishizaka, and T. Itaya, Tectonic setting of high-Mg andesite magmatism in SW Japan: K-Ar chronology of the Setouchi volcanic belt, *Geophys. J. Int.*, **144**, 625-631, 2001.
- Tatsumi, Y., and K. Sato, The western tip of the Setouchi Volcanic belt: constraints from K-Ar ages and geochemical characteristics of volcanic rocks from Oita Prefecture, *Hot spring Research of Oita Prefecture*, **52**, 15-19, 2001.
- Tatsumi, Y., T. Nakashima, and Y. Tamura, The petrology and geochemistry of calc-alkaline andesites on Shodo-Shima Island, SW Japan, *J. Petrol.*, **43**, 3-16, 2002.
- Tatsumi, Y., and T. Kogiso, The subduction factory: its role in the evolution of the Earth's crust and mantle, *Geol. Soc. London, Spec. Pub.*, in press, 2002.
- Tatsumi, Y., Some constraints on arc magma genesis, in *The Subduction Factory*, J. Eiler, ed., *Am. Geophys. Union, Monograph*, in press, 2002.
- Tsuboi, S., M. Saito, and Y. Ishihara, Verification of horizontal-to-vertical spectral ratio technique for estimation of site response using borehole seismographs, *Bull. Seismol. Soc. Am.*, **91**, 499-510, 2001.
- Tsuru, T., J. -O. Park, S. Miura, S. Kodaira, Y. Kido, and T. Hayashi, Along-arc structural variation of the plate boundary at the Japan Trench margin: implication of interplate coupling, *J. Geophys. Res.*, in press, 2002.
- Tsuru, T., J. -O. Park, N. Takahashi, S. Kodaira, Y. Kido, Y. Kaneda, and Y. Kono, Consideration on the updip limit of seismogenic zone of interplate earthquakes off Sanriku area from a seismic reflection data (in Japanese), *Butsuri Tansa*, **54**, 21-29, 2001.
- Tsuru, T., J. -O. Park, S. Miura, N. Takahashi, S. Kodaira, T. Higashikata, Y. Kido, and Y. Kaneda, Consideration on basal erosion along the plate boundary and interplate earthquake activity at the Japan Trench subduction zone, *Butsuri Tansa*, **54**, 21-29, 2001.
- Ujiie, K., Evolution and kinematics of an ancient decollement zone, melange in the Shimanto accretionary complex of Okinawa Island, Ryukyu Arc, *J. Struct. Geol.*, **24**, 937-952, 2002.
- Yamamoto, Y., K. Shimura, H. Tsunakawa, T. Kogiso, K. Uto, H. G. Barszczus, H. Oda, T. Yamazaki, and E. Kikawa, Geomagnetic paleosecular variation for the past 5 Ma in the Society Islands, French Polynesia, *Earth Planet Space*, **54**, 797-802, 2002.
- Yoshida, Y., K. Kanjo, S. Harada, K. Fujiwara, T. Ohtaki, A. Tanaka, S. Kaneshima, D. Suetsugu, S. Kamiya, Y. Ishihara, H. Takenaka, T. Saita, and Y. Fujii, Installation experiment of broadband seismometers in shallow vaults (in Japanese with English abstract), *Zisin*, **54**, 9-16, 2001.
- Wallace, P. J., F. A. Frey, D. Weis, and M. F. Coffin, Origin and evolution of the Kerguelen Plateau, Broken Ridge, and Kerguelen Archipelago: editorial, *J. Petrol.*, **43**, 1105-1108, 2002.
- Wei, Y. E., S. Yabuki, S. Kanayama, M. Honda, and Q. Chang, Geochemical characteristics of loess and paleosols in Yili Basin, NW China, *Proc. 10th Int. Symp. on Water Rock Interaction*, 1593-1596, 2001.
- Zheng, Z., H. Tanaka, Y. Tatsumi, and M. Kono, Basalt platforms in Inner Mongolia and Hebei Province, northeastern China: new K-Ar ages, geochemistries, and revision of paleomagnetic results, *Geophys. J. Int.*, **151**, 654-662, 2002.