

Eastern Indian Ocean Upwelling Research Initiative (EIOURI)


As IIOE-2/YMC overlapping interest

Weidong Yu, First Institute of Oceanography, SOA, China




With contributions from JAMSTEC, UT, WHOI, Scripps, UM, AMFR, LIPI, BMKG ...


Seasonal Upwelling Regions in Indian Ocean



SST (color) **Wind Velocity** in the Southwest Monsoon



Research Moored Array for African-Asian-Australian Monsoon Analysis and Prediction (RAMA)




Eastern Indian Ocean Upwelling Research Initiative (EIOURI)
Science Plan and Implementation Strategy Writing Group Meeting
9-12 Apr. 2014, Phuket, Thailand



Hosted by: Thailand-China Joint Laboratory for Climate and Marine Ecosystem

- Physical oceanography
 - Upwelling processes: local and remote forcing
 - Open ocean-coastal interactions: circulation and eddy
- BGC and Biological oceanography
 - Nutrient and plankton responses (e.g. classic vs cryptic upwelling, stoichiometry)
- Ecology
 - Food web, higher level responses and physical mechanisms (e.g. offshore advection, massive-kills)
- The human dimension

Pilot Cruise in 2013




Extension of EIOURI to YMC in collaboration with MOMSEI


- Indonesian-Australian Monsoon Onset
- In context of MOMSEI study in BoB
- To identify the first eastward propagating MJO which triggers Indonesian-Australian Monsoon

Monsoon Onset Process over BoB


(SST-MJO-Monsoon Onset)




MW + IR OI v3 Sea Surface Temperature: 2014/05/01 (~12 UTC) - Indian, North



Composite of BoB Monsoon Onset









FIG. 10. Composite plan view of OLR relative to onset at Darwin. Contour interval is 5 W m^{-2} with first contour at $\pm 10 \text{ W m}^{-2}$. The sequence is (a) day -8, (b) day +2, and (c) day +12.

Monsoon onset coincides with the first occurrence of convectively active 40-50 day oscillation.

Again, **SST-MJO-Onset** relationship should be examined, as in BoB.



Thanks!

