5.9 Ship-borne sky radiometer

(1) Personnel

Kazuma AOKI (University of Toyama) Tadahiro HAYASAKA (Tohoku University) Masataka SHIOBARA (NIPR)

- Principal Investigator / not onboard
- Co-Investigator / not onboard
- Co-Invectigator / not onboard

(2) Objectives

Objective of the observations in this aerosol is to study distribution and optical characteristics of marine aerosols by using a ship-borne sky radiometer (POM-01 MKII: PREDE Co. Ltd., Japan). Furthermore, collections of the data for calibration and validation to the remote sensing data were performed simultaneously.

(3) Methods and Instruments

Sky radiometer is measuring the direct solar irradiance and the solar aureole radiance distribution, has seven interference filters (0.34, 0.4, 0.5, 0.675, 0.87, 0.94, and 1.02 μ m). Analysis of these data is performed by SKYRAD.pack version 4.2 developed by Nakajima *et al.* (1996).

@ Measured parameters

- ① Aerosol optical thickness at five wavelengths (400, 500, 675, 870 and 1020 nm)
- 2 Ångström exponent
- ③ Single scattering albedo at five wavelengths
- (4) Size distribution of volume $(0.01 \,\mu\text{m} 20 \,\mu\text{m})$

GPS provides the position with longitude and latitude and heading direction of the vessel, and azimuth and elevation angle of sun. Horizon sensor provides rolling and pitching angles.

(4) Preliminary results

This study is not onboard. Data obtained in this cruise will be analyzed at University of Toyama.

(5) Data archives

Measurements of aerosol optical data are not archived so soon and developed, examined, arranged and finally provided as available data after certain duration. All data will archived at University of Toyama (K.Aoki, SKYNET/SKY: http://skyrad.sci.u-toyama.ac.jp/) after the quality check and submitted to JAMSTEC.

(6) Acknowledgment

The operations were supported by Global Ocean Development Inc.