

## <吉田尊雄業績目録>

(原著論文)

Kojima S., Kuriki Y., Yoshida T., Yazaki K., and Miura K.: "Fibril Formation by an Amphipathic  $\alpha$ -Helix-Forming Polypeptide Produced by Gene Engineering", Proc. Japan Acad., 73, Ser. B, 7-11 (1997).

Yoshida T., Yohda M., Iida T., Maruyama T., Taguchi H., Yazaki K., Ohta T., Odaka M., Endo I., and Kagawa Y.: "Structural and Functional Characterization of Homo-oligomeric Complexes of  $\alpha$  and  $\beta$  Chaperonin Subunits from the Hyperthermophilic Archaeum, *Thermococcus* strain KS-1", *J. Mol. Biol.*, **273** (3), 635-645 (1997).

Yoshida T., Yohda M., Iida T., Maruyama T., Taguchi H., Yazaki K., Ohta T., Odaka M., Endo I., and Kagawa Y.: [Corrigendum] "Structural and Functional Characterization of Homo-oligomeric Complexes of  $\alpha$  and  $\beta$  Chaperonin Subunits from the Hyperthermophilic Archaeum, *Thermococcus* strain KS-1", *J. Mol. Biol.*, **299** (5), 1399-1400 (2000).

Yoshida T., Wakiyama M., Yazaki K., and Miura K.: "Transmission electron and atomic force microscopic observation of polysomes on carbon-coated grids prepared by surface spreading", *J. Electron M.*, 46 (6), **503-506** (1997).

Yoshida, T., Yohda, M., Suzuki, M., Yazaki, K., Miura K., and Endo I. : "Characterization of Homo-oligomeric Complexes of  $\alpha$  and  $\beta$  Chaperonin Subunits from the Acidothermophilic Archaeon, *Sulfolobus* sp. strain 7", *Biochem. Biophys. Res. Comun.*, 242 (3), 640-647 (1998).

Furutani, M., Iida, T., Yoshida, T., Maruyama, T. :"Group II chaperonin in a thermophilic methanogen, *Methanococcus thermolithotrophicus*: chaperone activity and filament-forming ability", *J. Bio. Chem.*, 273 (43), 28399-28407 (1998).

Nojiri, M., Yohda, M., Odaka, M., Matsushita, Y., Tsujimura, M., Yoshida, T., Dohmae,

N., Takio, K., and Endo, I.: "Functional Expression of Nitrile Hydratase in Escherichia coli: Requirement of Nitrile Hydratase Activator and Post-translational Modification of Ligand Cysteine", *J. Biochem.*, 125 (4), 696-704 (1999).

Ideno, A., Yoshida, T., Furutani, M., and Maruyama, T.: "The 28.3kDa FK506 binding protein from a thermophilic archaeum, *Methanobacterium rmoautotrophicum*, protects the denatureation of proteins in vitro", *Eur. J. Biochem.*, 267 (11), 3139-3148 (2000).

Yazaki, K., Yoshida, T., Wakiyama, M., and Miura, K.: "Polysome of Eukaryotic Cells Observed by Electron Microscopy", *J. Electron M.*, 49 (5), 663-668 (2000)

Yoshida, T., Ideno, A., Hiyamuta, S., Yohda, M., and Maruyama, T.: "Natural chaperonin of the hyperthermophilic archaeum, *Thermococcus* strain KS-1; -A hetero-oligomeric chaperonin with variable subunit composition", *Mol. Microbiol.*, 39 (5), 1406-1413 (2001).

Ideno, A., Yoshida, T., Iida, T., Fuyutani, M., and Maruyama, T.: "FK506 binding protein of the hyperthermophilic archaea, *Thermococcus* sp. KS-1, a cold-shock inducible peptidyl-prolyl cis-trans isomerase with activites to trap and refold denatured proteins", *Biochemical. J.*, 357 (2), 465-471 (2001).

Usui, K., Yoshida, T., Maruyama, T., and Yohda, M.: "Small heat shock protein of a hyperthermophilic archaeum, *Thermococcus* sp. strain KS-1, exists as a spherical 24 mer and its expression is highly induced under heat-stress conditions", *J. Bioscience and Bioengineering*, 92 (2), 161-166 (2001).

Iizuka, R., Yoshida, T., Maruyama, T., Shomura, Y., Miki, K., and Yohda, M.: "Glycine at the 65th position plays an essential role in ATP dependent protein folding by archaeal group II chaperonin", *Biochem. Biophys. Res. Comum.*, 289 (5), 1118-1124 (2001).

Yoshida, T., Kawaguchi, R., Taguchi, H., Yoshida, M., Yasunaga, T., Wakabayashi, T., Yohda, M., and Maruyama, T.: "Archaeal Group II Chaperonin mediates protein folding

in the *cis*-cavity without a detachable GroES-like co-chaperonin", *J. Mol. Biol.*, 315 (1), 73-85 (2002).

Okouch, M., Yoshida, T., Maruyama, T., Kawarabayasi, Y., Kikuchi, H., and Yohda, M. :"*Pyrococcus* prefoldin stabilizes protein-folding intermediates and transfers them to chaperonins for correct folding", *Biochem. Biophys. Res. Comum.*, 291 (4), 769-779 (2002).

Yoshida, T., Kawaguchi, R., and Maruyama, T.: "Nucleotide specificity of an archaeal group II chaperonin from *Thermococcus* strain KS-1 with reference to the ATP-dependent protein folding cycle", *FEBS Lett.*, 514 (2-3), 269-274 (2002).

Yoshida, T., Ideno, A., Suzuki, R., Yohda, M., and Maruyama, T.: "Two kinds of archaeal group II chaperonin subunits with different thermostability in *Thermococcus* strain KS-1", *Mol. Microbiol.*, 44 (3), 761-769 (2002).

Shomura, Y., Yoshida, T., Maruyama, T., Yohda, M., and Miki, K.: "Crystallization and preliminary X-ray characterization of archaeal group II chaperonin alpha-subunit from *Thermococcus* strain KS-1", *Acta Crystallogr D Biol Crystallogr.*, 58 (Pt10 Pt2), 1830-1832 (2002).

Iizuka, R., Yoshida, T., Shomura, Y., Miki, K., Maruyama, T., Odaka, M., and Yohda, M.: "ATP binding is critical for the conformational change from an open to closed state in archaeal group II chaperonin", *J. Bio. Chem.*, 278 (45), 44959-44965 (2003).

Shomura, Y., Yoshida, T., Iizuka, R., Maruyama, T., Yohda, M., and Miki, K.: "Crystal structures of the group II chaperonin from *Thermococcus* strain KS-1: steric hindrance by the substituted amino acid, and inter-subunit rearrangement between two crystal forms", *J. Mol. Biol.*, 355 (5), 1265-1278 (2004).

Kohda, J., Yamada, T., Yoshida, T., Maruyama, T., Yohda, M., Fukuda, H., and Kondo, A.: "Refolding of proteins by hexadecamers and monomers of the a and b subunits of group II chaperonin from the hyperthermophilic archaeum *Thermococcus* strain KS-1,

*Bio. Eng. J.*, 18 (1), 73-79 (2004).

Iizuka, R., So, S., Inobe, T., Yoshida, T., Zako T., Kuwajima, K., and Yohda, M.: "Role of the helical protrusion in the conformational change and molecular chaperone activity of the archaeal group II chaperonin", *J. Bio. Chem.*, 279 (18), 18834-18839 (2004).

Furutani, M., Hata, J., Shomura, Y., Itami, K., Yoshida, T., Izumoto, Y., Togi, A., Ideno, A., Yasunaga, T., Miki, K., and Maruyama, T.: "An engineered chaperonin caging a guest protein: Structural insights and potential as a protein expression tool", *Protein Sci.*, 14 (2), 341-350 (2005).

Iizuka, R., Yoshida, T., Ishii, N., Zako, T., Takahashi, K., Maki, K., Inobe, T., Kuwajima, K., and Yohda, M.: "Characterization of archaeal group II chaperonin-ADP-metal fluoride complexes: implications that group II chaperonins operate as a "two-stroke engine""", *J. Bio. Chem.*, 280 (48), 40375-40383 (2005).

Yoshida, T., Kanzaki, T., Iizuka, R., Komada, T., Zako, T., Suzuki, R., Maruyama, T. and Yohda, M. : "Contribution of the C-terminal region to the thermostability of the archaeal group II chaperonin from *Thermococcus* sp. strain KS-1 ", *Extremophiles*, 10 (5), 451-459 (2006).

Zako, T., Murase, Y., Iizuka, R., Yoshida, T., Kanzaki, T., Ide, N., Maeda, M., Funatsu, T. and Yohda, M.: "Localization of Prefoldin Interaction Sites in the Hyperthermophilic Group II Chaperonin and Correlations between Binding Rate and Protein Transfer Rate", *J. Mol. Biol.*, 364(1), 110-120 (2006).

Yoshida, T., Iizuka, R., Itami, K., Yasunaga, T., Sakuraba, H., Ohshima, T., Yohda, M. and Maruyama, T.: "Comparative analysis of the protein folding activities of two chaperonin subunits of *Thermococcus* strain KS-1; the effects of beryllium fluoride", *Extremophiles*, 11(2), 225-235 (2007).

Kuwahara, H., Yoshida, T., Takaki, Y., Shimamura, S., Nishi, S., Harada, m.,

Matsuyama, K., Takishita, K., Kawato, M., Uematsu, K., Fujiwara, Y., Sato, T., Kato, C., Kitagawa, M., Kato, I., and Maruyama T.: "Reduced genome of the thioautotrophic intracellular symbiont in a deep-sea clam, *Calyptogena okutanii*", *Curr biol* 17(10), 881-886 (2007).

Iizuka, R., Sugano, Y., Ide, N., Ohtaki, A., Yoshida, T., Fujiwara, S., Imanaka, T., and Yohda, M.: "Functional characterization of recombinant prefoldin complexes from a hyperthermophilic archaeon, *Thermococcus* sp. strain KS-1", *J. Mol. Biol.* 377(3), 972-983 (2008).

Saji, H., Iizuka, R., Yoshida, T., Abe, T., Kidokoro, S., Ishii, N., and Yohda, M.: "Role of IXI/V motif in oligomer assembly and function of StHsp14.0, a small heat shock protein from the acidothermophilic archaeon, *Sulfolobus tokodaii* strain 7", *PROTEINS: Structure, Function, and Bioinformatics*, 71(2), 771-782 (2008).

Kuwahara, H., Takaki, Y., Yoshida, T., Shimamura, S., Takishita, K., Reimer, J.D., Kato, C., and Maruyama T.: "Reductive genome evolution in chemoautotrophic intracellular symbionts of deep-sea *Calyptogena* clams", *Extremophiles*, 12(3), 365-374 (2008).

Fujikura K., Sasaki T., Yamanaka T., and Yoshida T. Turrids whelk, *Phymorhynchus buccinoides* feeds on Bathymodiolus mussels at a seep site in Sagami Bay, Japan. Plankton and Benthos Research 4(1), 23-30 (2009).

Harada M., Yoshida T., Kuwahara H., Shimamura S., Takaki Y., Kato C., Miwa T., Miyake H., and Maruyama T.: "Expression of genes for sulfur oxidation in the intracellular chemoautotrophic symbiont of the deep-sea bivalve *Calyptogena okutanii*", *Extremophiles* 13(6), 895-903 (2009).

Takishita K., Kakizoe N., Yoshida T., and Maruyama T.: "Molecular evidence that phylogenetically diverged ciliates are active in microbial mats of Deep-Sea Cold Seep Sediment", *J Eukaryot Microbiol.* 57(1), 76-86 (2010).

Koito T., Nakamura-Kusakabe I., Yoshida T., Mauyama T., Omata T., Miyazaki N., and Inoue K.: "Effect of long-term exposure to sulfides on taurine transporter gene expression in the gill of the deep-sea mussel *Bathymodiolus platifrons*, which harbors a methanotrophic symbiont" *Fish. Sci.* 76(2), 381-388 (2010).

Koito, T., Morimoto, S., Toyohara H., Yoshida, T., Jimbo, M., Maruyama, T., Miyazaki, N., Inoue, K.: Decline in taurine transporter mRNA and thioautotrophic bacterial 16S rDNA levels after transplantation of the hydrothermal-vent mussel *Bathymodiolus septemdierum* to a non-vent position. *Cah. Biol. Mar.* 51, 429-433 (2010).

Kuwahara H., Takaki, Y., Shimamura, S., Yoshida, T., Maeda, T., Kunieda, T., and Maruyama, T. : Loss genes for DNA recombination and repair in the reductive genome evolution of thioautotrophic symbionts of Calyptogena clams. *BMC Evol. Biol.* 11(1), 285 (2011).

Maeda T, Hirose E, Chikaraishi Y, Kawato M, Takishita K, Yoshida, T., Verbruggen, H., Tanaka, J., Shimamura, S., Takaki, Y., Tsuchiya, M., Iwai, K., and Maruyama, T. (2012) Algivore or Phototroph? *Plakobranchus ocellatus* (Gastropoda) Continuously Acquires Kleptoplasts and Nutrition from Multiple Algal Species in Nature. *PLoS ONE* 7(7): e42024. doi:10.1371/journal.pone.0042024

Uchida, H., Hamana, K., Miyazaki, M., Yoshida, T., and Nogi, Y. : *Parasphingopyxis lamellibrachiae* gen. nov., sp. nov., isolated from a marine annelid worm. *IJSEM* 62, 2224-2228 (2012).

Noguchi, F., Kawato, M., Yoshida, T., Fujiwara, Y., Fujikura, K., and Takishita, K.:”A Novel Alveolate in Bivalves with Chemosynthetic Bacteria Inhabiting Deep-Sea Methane Seeps”, Journal of Eukaryotic Microbiology 60, 158-165 (2013).

Hori, S., Tsuchiya, M., Nishi, S., Arai, W., Yoshida, T., and Takami, H. :”Active bacterial flora surrounding foraminifera (xenophyophorea) living on the deep-sea floor”, Bioscience, Biotechnology, and Biochemistry 77, 381-384 (2013).

Watanabe, H., Seo, E., Takahashi, Y., Yoshida, T., Kojima, S., Fujikura, K., and Miyake, H. :” Spatial distribution of sister species of vesicomyid bivalves *Calyptogena okutanii* and *Calyptogena soyoae* along an environmental gradient, in chemosynthetic biological communities in Japan”, Journal of Oceanography 69(1), 129-134 (2013).

Nakamura, Y., Konishi, M., Ohishi, K., Kusaka, C., Tame, A., Hatada, Y., Fujikura, K., Nakazawa, M., Fujishima, M., Yoshida, T., and Maruyama, T. :”Mucus glycoproteins selectively secreted from bacteriocytes in gill filaments of the deep-sea clam *Calyptogena okutanii”*, Open Journal of Marine Science, 3(4), 167-174 (2013).

Hongo, Y., Nakamura, Y., Shimamura, S., Takaki, Y., Uematsu, K., Toyofuku, T., Hirayama, H., Takai, K., Nakazawa, M., Maruyama, T., and Yoshida, T. :”Exclusive localization of carbonic anhydrase in bacteriocytes of the deep-sea clam *Calyptogena okutanii* with thioautotrophic symbiotic bacteria”, Journal of Experimental Biology, 216(23), 4403-4414 (2013).

Ohishi, K., Suzuki, R., Maeda, T., Tsuda, M., Abe, E., Yoshida, T., Endo, Y., Okamura, M., Nagamine, T., Yamamoto, H., Ueda, M., and Maruyama, T. :”Recent host range expansion of canine distemper virus and variation in its receptor, the signaling lymphocyte activation molecule, in carnivores”, Journal of Wildlife Diseases, 50(3), 596-606 (2014).

Numoto, N., Nakagawa, T., Ohara, R., Hasegawa, T., Kita, A., Yoshida, T., Maruyama, T., Imai, K., Fukumori, Y., and Miki, K. :”The structure of a deoxygenated 400 kDa

hemoglobin reveals ternary and quaternary structural changes of giant hemoglobins”,  
*Acta Crystallogr D Biol Crystallogr.*, **70**, 1823-1831 (2014).

Takihara, H., Ogiara, J., Yoshida, T., Okuda, S., Nakajima, M., Iwabuchi, N., and Sunairi, M. :” Enhanced translocation and growth of *Rhodococcus erythropolis* PR4 in the alkane phase of aqueous-alkane two phase cultures were mediated by GroEL2 overexpression”, *Microbes and Environments*, **29**, 346-352 (2014).

Shigeno, S., Ogura, A., Mori, T., Toyohara, H., Yoshida, T., Tsuchida, S., and Fujikura, K. : ” Sensing deep extreme environments: the receptor cell types, brain centers, and multi-layer neural packaging of hydrothermal vent endemic worms” , *Frontiers in Zoology*, 11:82 (2014).

Nagasaki, T., Hongo, Y., Koito, Y., Nakamura-Kusakabe, I., Shimamura, S., Takaki, Y., Yoshida, T., Maruyama, T., and Inoue, K.: ” Cysteine dioxygenase and cysteine sulfinate decarboxylase genes of the deep-sea mussel *Bathymodiolus sepiemdierum*: possible involvement in hypotaurine synthesis and adaptation to hydrogen sulfide” , *Amino Acids*, 47, 571–578 (2015).

Tame, A., Yoshida, T., Ohishi, K., and Maruyama, T.:”Phagocytic activities of hemocytes from the deep-sea symbiotic mussels *Bathymodiolus japonicus*, *B. platifrons*, and *B. sepiemdierum*.”, *Fish and Shellfish Immunology*, **45**, 146-156 (2015).

Sato, H., Nakasone, K., Yoshida, T., Kato, C., and Maruyama,T.:”Increases of heat-shock proteins and their mRNAs at high hydrostatic-pressure in a deep-sea piezophilic bacterium, *Shewanella violacea*”, *Extremophiles*, in press

(プロシードィングス)

Yohda, M., Honda, J., Nagamune, T.. Endo, I., Yoshida, T., and Miura, K. : "Molecular cloning and nucleotide sequence of the gene coding photosensitive nitrile hydratase", *Ann N Y Acad Sci* 721, 158-159 (1994).

Koito, T., Yoshida, T., Maruyama, T., Miyazaki, N., Inoue, K. :"Quantitative analyses of taurine transporter mRNA in hydrothermal vent and cold seep mussels *Bathymodiolus septemfasciatus* and *B. platifrons*", 5th World Fisheries Congress (2009).

### (総説)

養王田正文, 吉田尊雄, 遠藤勲：“古細菌のシャペロニン”, バイオサイエンスとインダストリー, 55巻, No. 9, 623-626 (1997)

吉田尊雄, 養王田正文, 丸山正：“古細菌の分子シャペロンの世界 シャペロニンを中心にして”, 蛋白質核酸酵素, 48巻, No. 1, 33-39 (2003)

吉田尊雄, 碓井啓資, 飯塚怜, 座古保, 養王田正文：“古細菌の分子シャペロンシステムの反応機構”, 蛋白質核酸酵素, 49巻, No. 7, 858-861 (2004)

養王田正文, 飯塚怜, 座古保, 吉田尊雄：“超好熱性古細菌の分子シャペロン”, 酵素工学ニュース, 第52号, 6-12 (2004)

Iizuka, R., So, S., Inobe, T., Yoshida, T., Zako, T., Kuwajima, K, Yohda, M.: "Role of the helical protrusion in the conformational change and molecular chaperone activity of the archaeal group II chaperonin", Photon Factory Activity Report, 22, 48-49, (2005).

吉田尊雄、丸山正：“ストレスタンパク質を利用した海洋微生物の現場計測の可能性”, 月刊海洋, 第38号, No. 12, 837-843 (2006)

### (本)

吉田尊雄：“熱水にふれてもなぜ生きていられる！？”, 潜水調査船が観た深海生物 深海生物研究の現在, 藤倉克則、奥谷喬司、丸山正、164-165 (2008)

丸山正、藤原義弘、吉田尊雄：“深海における海産無脊椎動物と微生物の共生-化学合成共生系を中心にして”, 潜水調査船が観た深海生物 深海生物研究の現在, 藤倉克則、奥谷喬司、丸山正、37-50 (2008)

吉田尊雄、高木善弘、島村繁、丸山正 化学合成生態系の無脊椎動物- 微生物  
間細胞内共生系からみた共生菌のゲノム縮小進化 「難培養微生物研究の最新  
技術 II」、CMC出版、126-135(2010)

丸山正、高木善弘、吉田尊雄：“深海化学合成共生細菌のゲノムと共生の進化”，  
生物の科学 遺伝、65 (6) , 35-43 (2011)

**Jimbo, M., S. Satoh, H. Hasegawa, H. Miyake, T. Yoshida, T. Maruyama,  
and H. Kamiya** (2012) The difference of lectin recovery by sugar-bound  
resin, Affinity Chromatography Dr. Sameh Magdeldin (Ed.), ISBN:  
978-953-51-0325-7, InTech, Available from:  
[http://www.intechopen.com/books/affinity-chromatography/the-difference  
-of-lectin-recovery-by-sugar-boundresins](http://www.intechopen.com/books/affinity-chromatography/the-difference-of-lectin-recovery-by-sugar-boundresins)

高木善弘、吉田尊雄：“化学合成生態系における動物と化学合成細菌との共生”，  
生物の科学 遺伝、67 (4) , 482-489 (2013)